

ON ORBITAL VARIETY CLOSURES IN $\mathfrak{sl}_n$

II. DESCENDANTS OF A RICHARDSON ORBITAL VARIETY

ANNA MELNIKOV

ABSTRACT. For a semisimple Lie algebra $\mathfrak{g}$ the orbit method attempts to assign representations of $\mathfrak{g}$ to (coadjoint) orbits in $\mathfrak{g}^*$. Orbital varieties are particular Lagrangian subvarieties of such orbits leading to highest weight representations of $\mathfrak{g}$. In $\mathfrak{sl}_n$ orbital varieties are described by Young tableaux. In this paper we consider so called Richardson orbital varieties in $\mathfrak{sl}_n$. A Richardson orbital variety is an orbital variety whose closure is a standard nilradical. We show that in $\mathfrak{sl}_n$ a Richardson orbital variety closure is a union of orbital varieties. We give a complete combinatorial description of such closures in terms of Young tableaux.

This is the second paper in the series of three papers devoted to a combinatorial description of orbital variety closures in $\mathfrak{sl}_n$ in terms of Young tableaux.

1. Introduction

1.1. This is a continuation of Part I ([3]) whose notation we retain. Our main goal in this series of papers is to give a description of an orbital variety closure in $\mathfrak{sl}_n$. In this paper we give a complete description of the closure of a so called Richardson orbital variety.

For the convenience of the reader we repeat necessary notation and results from Part I which can be formulated in short. If the formulation is too long, as for example, in the case of Robinson-Schensted procedure we provide the exact reference to the subsection of Part I.

1.2. Let $\mathbf{G}$ be a connected semisimple finite dimensional complex algebraic group. Let $\mathfrak{g}$ be its Lie algebra. Fix some triangular decomposition $\mathfrak{g} = \mathfrak{n} \oplus \mathfrak{h} \oplus \mathfrak{n}^-$. Take some $x \in \mathfrak{n}$ and consider $\mathcal{O} = \mathcal{O}_x$ a nilpotent orbit of $\mathfrak{g}$ under the adjoint action of $\mathbf{G}$ (that is $\mathcal{O} = \mathbf{G}x = \{gxg^{-1} \mid g \in \mathbf{G}\}$). Since $\mathfrak{g}$ is semi-simple we can identify $\mathfrak{g}^*$ with $\mathfrak{g}$ through the Killing form. This identification gives an adjoint orbit a symplectic structure. By the results of R. Steinberg, N. Spaltenstein and A. Joseph (cf. [1]) $\mathcal{O} \cap \mathfrak{n}$ is equidimensional and Lagrangian. Its irreducible components are called orbital varieties associated to $\mathcal{O}$.

Given an orbital variety $\mathcal{V}$ we denote by $\mathcal{O}_{\mathcal{V}}$ the orbit $\mathcal{V}$ is associated to.

This work was partially supported by the EEC program TMR-grant ERB FMRX-C T97-0100.

1.3. Let $R \subset \mathfrak{h}^*$ denote the set of non-zero roots, R^+ the set of positive roots corresponding to $\mathfrak{n}$ and $\Pi \subset R^+$ the resulting set of simple roots. Let $X_\alpha := \mathbb{C}x_\alpha$ denote the root subspace corresponding to $\alpha \in R$. Then $\mathfrak{n} = \bigoplus_{\alpha \in R^+} X_\alpha$. Set $\mathfrak{n} \cap^w \mathfrak{n} := \bigoplus_{\alpha \in R^+ \cap^w R^+} X_\alpha$.

Let $\mathbf{B}$ be the standard Borel subgroup of $\mathbf{G}$, i.e. such that $\text{Lie}(\mathbf{B}) = \mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}$. $\mathbf{B}$ acts by conjugation on $\mathfrak{n}$ and its subsets. Let W be the Weyl group for the pair $(\mathfrak{g}, \mathfrak{h})$. By Steinberg's construction there exists a surjection from W onto the set of orbital varieties defined by $w \rightarrow \overline{\mathbf{B}(\mathfrak{n} \cap^w \mathfrak{n})} =: \overline{\mathcal{V}}_w$ (cf. Part I, 2.1.2, 2.1.3).

Note that there exists the unique nilpotent orbit $\mathcal{O}_w$ such that $\overline{\mathcal{O}}_w = \overline{\mathbf{G}(\mathfrak{n} \cap^w \mathfrak{n})}$. Obviously one has $\mathcal{O}_w = \mathcal{O}_{\mathcal{V}_w}$.

1.4. Take $\mathcal{I} \subset \Pi$, let $\mathbf{P}_{\mathcal{I}}$ denote the unique standard parabolic subgroup of $\mathbf{G}$ generated by the standard parabolic subgroups $\mathbf{P}_\alpha : \alpha \in \mathcal{I}$. Let $\mathbf{M}_{\mathcal{I}}$ be the unipotent radical of $\mathbf{P}_{\mathcal{I}}$ and $\mathfrak{m}_{\mathcal{I}} = \text{Lie}(\mathbf{M}_{\mathcal{I}})$ be the corresponding nilradical in $\mathfrak{n}$.

Let $W_{\mathcal{I}} := \langle s_\alpha : \alpha \in \mathcal{I} \rangle$ be the corresponding parabolic subgroup of W . Let $w_{\mathcal{I}}$ be the unique longest element of $W_{\mathcal{I}}$, that is such that

$$w_{\mathcal{I}}(\alpha) \in \begin{cases} R^- & \text{if } \alpha \in \mathcal{I} \\ R^+ & \text{if } \alpha \in R^+ \setminus \sum_{\beta \in \mathcal{I}} \mathbb{N} \cdot \beta \end{cases} \quad (*)$$

Note that $\mathfrak{n} \cap^{w_{\mathcal{I}}} \mathfrak{n} = \mathfrak{m}_{\mathcal{I}}$. Thus $\overline{\mathcal{V}}_{w_{\mathcal{I}}} = \overline{\mathbf{B}(\mathfrak{n} \cap^{w_{\mathcal{I}}} \mathfrak{n})} = \mathfrak{m}_{\mathcal{I}}$. Just to simplify the notation we denote $\mathcal{V}_{\mathcal{I}} := \mathcal{V}_{w_{\mathcal{I}}}$.

Set $\mathcal{O}_{\mathcal{I}} := \mathcal{O}_{\mathcal{V}_{\mathcal{I}}}$. One calls $\mathcal{V}_{\mathcal{I}}$ a Richardson orbital variety (associated to $\mathcal{O}_{\mathcal{I}}$) defined by $\mathcal{I}$.

1.5. From now on we consider only the case of $\mathfrak{g} = \mathfrak{sl}_n$. Every nilpotent orbit in $\mathfrak{g}$ is defined by its Jordan form which in turn is completely defined by the set of lengths of its Jordan blocks. So we get a bijection φ from the set of partitions of n onto the set of nilpotent orbits in $\mathfrak{g}$. We write a partition in decreasing order that is $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ where $\sum_{i=1}^k \lambda_i = n$ and $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_k > 0$. We set $\mathcal{O}_\lambda := \varphi(\lambda)$.

Define a partial order on partitions of n as follows. Let $\lambda = (\lambda_1, \dots, \lambda_k)$ and $\mu = (\mu_1, \dots, \mu_j)$ be partitions of n . If $j \neq k$ complete the partition with less number of elements by 0 so that we can consider that both partitions have $\max(j, k)$ elements. Define $\lambda \leq \mu$ if for each $i : 1 \leq i \leq \max(j, k)$ one has

$$\sum_{m=1}^i \lambda_m \geq \sum_{m=1}^i \mu_m.$$

Then by the result of Gerstenhaber (cf. Part I 2.3.2)

$$\overline{\mathcal{O}}_\lambda = \coprod_{\mu \geq \lambda} \mathcal{O}_\mu.$$

We define a partial order on nilpotent orbits by $\mathcal{O} \leq \mathcal{O}'$ if $\mathcal{O}' \subset \overline{\mathcal{O}}$. We say that $\mathcal{O}' : \overline{\mathcal{O}'} \subsetneq \overline{\mathcal{O}}$ is a descendant of $\mathcal{O}$ if for any $\mathcal{O}''$ such that $\mathcal{O} \leq \mathcal{O}'' \leq \mathcal{O}'$ one has $\mathcal{O}'' = \mathcal{O}$ or $\mathcal{O}'' = \mathcal{O}'$.

1.6. Recall the notation and the conventions of Part I, 2.2.1. In particular recall that in our case $\Pi = \{\alpha_i\}_{i=1}^{n-1}$. Recall from Part I, 2.4.5, 2.4.6 the combinatorial characterization of orbital varieties in $\mathfrak{g}$ in terms of Young tableaux. It is constructed as follows. As we explained in 1.3 by Steinberg construction one has a surjection from Weyl group W onto the set of orbital varieties. Set $\mathcal{C}_{\mathcal{V}} := \{w \in W \mid \overline{\mathcal{V}} = \overline{\mathcal{V}}_w\}$. We get a partition of Weyl group into so called geometric cells. We also define $\mathcal{C}_w := \mathcal{C}_{\mathcal{V}_w}$.

Identify Weyl group of $\mathfrak{sl}_n$ with symmetric group $\mathbf{S}_n$ (cf. Part I, 2.2.2). Robinson-Schensted procedure (cf. Part I, 2.4.6) gives a surjection $w \mapsto T(w)$ from $\mathbf{S}_n$ onto $\mathbf{T}_n$ the set of standard Young tableaux with n entries. By R. Steinberg (cf. Part I, 2.4.6) one has $y \in \mathcal{C}_w$ if and only if $T(y) = T(w)$. In such a way we get a bijection $\phi(T(w)) := \mathcal{V}_w$ from the set of standard Young tableaux onto the set of orbital varieties. Given a standard Young tableau T set $\mathcal{V}_T := \phi(T)$. Respectively given an orbital variety $\mathcal{V}$ set $T_{\mathcal{V}} := \phi^{-1}(\mathcal{V})$.

Given a Young tableau T let $\text{sh}(T) = \lambda$ be the partition of n from which T was built, that is $\lambda = (\lambda_1, \dots, \lambda_k)$ where λ_i is the length of i -th row of T . Then $\mathcal{V}_T$ is associated to $\mathcal{O}_{\lambda}$ so that the characterization of orbital varieties by Young tableaux is compatible with the characterization of nilpotent orbits by partitions.

1.7. We give a combinatorial description of a Richardson orbital variety closure in terms of Young tableaux in the spirit of Gerstenhaber's construction.

Given a standard Young tableau $T \in \mathbf{T}_n$ and a natural number $u : 1 \leq u \leq n$ let $r_T(u)$ denote the number of the row of T u belongs to. Recall from Part I, 2.4.14 that $\tau(T) = \{\alpha_i : r_T(i+1) > r_T(i)\}$. Recall also from there that $\tau(\mathcal{V}_T) = \tau(T)$.

We show in 2.3 that

$$\overline{\mathcal{V}}_T = \bigcup_{\tau(\mathcal{V}) \supseteq \tau(T)} \mathcal{V}. \quad (**)$$

This formula resembles Gerstenhaber's result in its simplicity and clearness.

1.8. We define a partial order on the set of orbital varieties by $\mathcal{V} \stackrel{G}{\leq} \mathcal{W}$ if $\mathcal{W} \subset \overline{\mathcal{V}}$. We call it the geometric order. This order is compatible with the order on nilpotent orbits defined in 1.5 in the sense that $\mathcal{V} \stackrel{G}{\leq} \mathcal{W}$ implies that $\mathcal{O}_{\mathcal{V}} \leq \mathcal{O}_{\mathcal{W}}$. Moreover by Part I, 4.1.8 for any orbital variety $\mathcal{V}$ in $\mathfrak{sl}_n$ and any nilpotent orbit $\mathcal{O}$ such that $\mathcal{O} > \mathcal{O}_{\mathcal{V}}$ there exist an orbital variety $\mathcal{W}$ associated to $\mathcal{O}$ such that $\mathcal{V} \stackrel{G}{<} \mathcal{W}$.

We say that $\mathcal{W} : \mathcal{W} \stackrel{G}{>} \mathcal{V}$ is a geometric descendant of $\mathcal{V}$ if for any $\mathcal{Y}$ such that $\mathcal{V} \stackrel{G}{\leq} \mathcal{Y} \stackrel{G}{\leq} \mathcal{W}$ one has $\mathcal{Y} = \mathcal{V}$ or $\mathcal{Y} = \mathcal{W}$.

The description of the set of descendants of a nilpotent orbit is obtained as an easy corollary of Gerstenhaber's formula (cf. Part I, 2.3.3). However the description of the set of geometric descendants of a Richardson orbital variety cannot be easily obtained from 1.7 (**). Yet the result is very clear-cut. This is the main result of the paper. Its formulation demands some additional combinatorial notation so we refer the reader to 2.8 for the exact statement of the result.

1.9. By [5] $\mathfrak{n} \cap \mathcal{O}$ is a union of orbital varieties associated to $\mathcal{O}$. Moreover by [4]

$$\overline{\mathfrak{n} \cap \mathcal{O}} = \mathfrak{n} \cap \overline{\mathcal{O}}.$$

As we show in 2.3 for any $\mathcal{O} \subset \overline{\mathcal{O}}_{\mathcal{I}}$ one has $\mathfrak{m}_{\mathcal{I}} \cap \mathcal{O}$ is a union of all orbital varieties (associated to $\mathcal{O}$) whose τ -invariant contains $\mathcal{I}$. A natural question is whether for such $\mathcal{O}$ one has as well that

$$\overline{\mathfrak{m}_{\mathcal{I}} \cap \mathcal{O}} = \mathfrak{m}_{\mathcal{I}} \cap \overline{\mathcal{O}}. \quad (***)$$

Theorem 2.8 provides in particular that $\mathcal{V}$ being a descendant of $\mathcal{V}_{\mathcal{I}}$ does not necessarily imply that $\mathcal{O}_{\mathcal{V}}$ is a descendant of $\mathcal{O}_{\mathcal{I}}$. Using this fact we show in 4.1 that in general equality (***) does not hold.

1.10. In [2] the ideals of definition of an orbital variety closure of codimension 1 in some nilradical were constructed as well as strong quantization of such orbital variety. The description of the set of geometric descendants of a Richardson orbital variety is central for the generalization of the results of [2] to orbital varieties of codimension greater than 1 in a nilradical.

1.11. A natural and interesting problem is to determine which orbital variety closures are complete intersections. This is very important in particular for strong quantization of an orbital variety. It is more or less obvious that the majority of orbital varieties are not complete intersections. Concentrating on the descendants of a Richardson orbital variety one can see at once that all the descendants of codimension 1 in $\mathfrak{m}_{\mathcal{I}}$ are complete intersections just by Krull theorem. What can be said about descendants of a Richardson orbital variety of codimension greater than 1 in $\mathfrak{m}_{\mathcal{I}}$? It is very easy to check that in $\mathfrak{sl}_n$ for $n \leq 5$ all the descendants of a Richardson orbital variety are complete intersections. We give an example in $\mathfrak{sl}_6$ of $\mathcal{V}_{\mathcal{I}}$ and its descendant $\mathcal{W}$ such that

- (i) $\text{codim}_{\mathfrak{m}_{\mathcal{I}}} \overline{\mathcal{W}} = 2$;
- (ii) $\mathcal{O}_{\mathcal{W}}$ is a descendant of $\mathcal{O}_{\mathcal{I}}$;
- (iii) $\mathcal{W}$ is not a complete intersection.

We construct the ideal of definition for this $\mathcal{W}$.

This example shows that even among the descendants of a Richardson orbital variety the majority of those of codimension greater than 1 in $\mathfrak{m}_{\mathcal{I}}$ are not complete intersections.

1.12. The body of the paper consists of three sections.

In Section 2 we develop the combinatoric notation essential to state the main theorem (formulated in 2.8).

Section 3 is devoted to the proof of theorem 2.8. Here we develop the combinatorics connected to the non-trivial involution ψ of the Dynkin diagram of $\mathfrak{sl}_n$. This involution induces involution on Weyl group W . Given $w \in W$ in word form we give a formula for $\psi(w)$ in a word form in 3.8. As well involution ψ induces order preserving involution on the set of Young tableaux. In general there is no straightforward formula determining $\psi(T)$. However we develop such formulas for the tableaux considered in the paper and use them as one of the central tools in our proof.

Finally in section 4 we discuss in detail 1.9 and 1.11.

In the end one can find the index of notation in which symbols appearing frequently are given with the subsection where they are defined.

2. A Richardson orbital variety closure and its descendants

2.1. Let us recall the notation and the results concerning Weyl group and in particular $\mathbf{S}_n$ from Part I that we use in this section.

For $w \in W$ set $S(w) := R^+ \cap^w R^-$. The map $w \mapsto S(w)$ is injective, so we define a partial order relation on W by taking $y \leq^D w$ if $S(y) \subseteq S(w)$. It is called the Duflo order.

Recall notation X_α from 1.3. Since $\mathfrak{n} \cap^w \mathfrak{n} = \bigoplus_{\alpha \in R^+ \cap^w R^+} X_\alpha$ one has $y \leq^D w$ if and only if $\mathfrak{n} \cap^w \mathfrak{n} \subseteq \mathfrak{n} \cap^y \mathfrak{n}$. In particular if $y \leq^D w$ then $\mathcal{V}_w \leq^G \mathcal{V}_y$.

Set $\tau(w) := S(w) \cap \Pi$. Recall notion of the standard Borel subalgebra $\mathfrak{b}$ from 1.3. For $\alpha \in \Pi$ let $\mathbf{P}_\alpha$ be the standard parabolic subgroup such that $\text{Lie}(\mathbf{P}_\alpha) = \mathfrak{b} \oplus X_{-\alpha}$. Given an orbital variety $\mathcal{V}$ set $\mathbf{P}_\alpha(\mathcal{V}) := \{gXg^{-1} \mid X \in \mathcal{V}, g \in \mathbf{P}_\alpha\}$ and let $\tau(\mathcal{V}) := \{\alpha \in \Pi \mid \mathbf{P}_\alpha(\mathcal{V}) = \mathcal{V}\}$.

Let us consider $\mathfrak{g} = \mathfrak{sl}_n$. In that case $\Pi = \{\alpha_i\}_{i=1}^{n-1}$ and any $\alpha \in R^+$ is $\alpha_{i,j} = \sum_{k=i}^{j-1} \alpha_k$ for some $i, j : 1 \leq i < j \leq n$. As well in that case W is identified with $\mathbf{S}_n$ (cf. Part I, 2.2.1, 2.2.2). We write $w \in \mathbf{S}_n$ in a word form $w = [a_1, \dots, a_n]$ where $a_i = w(i)$. Put $p_w(i) = j$ if $i = a_j$, that is $p_w(i)$ is its place (index) in word w . By Part I, 2.2.4 one has $\alpha_{i,j} \in S(w)$ iff $p_w(i) > p_w(j)$.

Recall $\tau(T)$ from 1.7. By Part I, 2.1.7, 2.4.14 $\tau(w) = \tau(\mathcal{V}_w) = \tau(T(w))$.

We also need the following notation from Part I, 2.2.5. Given a word $w = [a_1, \dots, a_n]$ let $<w> := \{a_i\}_{i=1}^n$ be the set of its entries.

- (i) Set $\bar{w} := [a_n, \dots, a_1]$ to be the word with order reverse to the order of w .
- (ii) Given words $x = [a_1, \dots, a_k]$, $y = [b_1, \dots, b_m]$ such that $\langle x \rangle \cap \langle y \rangle = \emptyset$ set $[x, y] = [a_1, \dots, a_k, b_1, \dots, b_m]$ to be a colligation.

Given a fixed set E of n distinct positive integers we let $\mathbb{S}_n$ or $\mathbb{S}_E$ denote the set of words w such that the set of its entries $\langle w \rangle = E$.

2.2. As well we need to recall some notation concerning Young tableaux. Given a Young tableau T recall from Part I, 2.4.2 that T_j^i is the entry on the intersection of i -th row and j -th column and $\omega^i(T)$ is the last entry of row i of T . As well T_j is j -th column of T ; T^i is i -th row of T ; notation $T^{i,j}$ means subtableau of T containing all the rows from i -th row to j -th row and finally $T^{i,\infty}$ is the subtableau of T containing all the rows from i -th row and down. Put $|T_j|$ (resp. $|T^i|$) to be the length of j -th column (resp. i -th row) of T .

For example if

$$T = \begin{array}{|c|c|c|c|c|} \hline 1 & 2 & 4 & 8 & 9 \\ \hline 3 & 5 & 6 & & \\ \hline 7 & & & & \\ \hline 10 & & & & \\ \hline \end{array}$$

then $T_2^2 = 5$, $T_4^1 = 8$; $\omega^1(T) = 9$, $\omega^2(T) = 6$, $\omega^3(T) = 7$, $\omega^4(T) = 10$ and

$$T_3 = \begin{array}{|c|} \hline 4 \\ \hline 6 \\ \hline \end{array}, \quad T^2 = \begin{array}{|c|c|c|} \hline 3 & 5 & 6 \\ \hline \end{array}, \quad T^{2,3} = \begin{array}{|c|c|c|} \hline 3 & 5 & 6 \\ \hline 7 & & \\ \hline \end{array}, \quad T^{2,\infty} = \begin{array}{|c|c|c|} \hline 3 & 5 & 6 \\ \hline 7 & & \\ \hline 10 & & \\ \hline \end{array}$$

Given a tableau T let $\langle T \rangle$ denote the set of its entries. Given a fixed set E of n distinct positive integers, let $\mathbb{T}_n$ or $\mathbb{T}_E$ be the set of Young tableaux T such that $\langle T \rangle = E$.

Recall from Part I, 2.4.3 the definition of a tableau (T, S) where $\langle T \rangle \cap \langle S \rangle = \emptyset$. Note that if for every row $i : 1 \leq i \leq \min(|S_1|, |T_1|)$ one has $\omega^i(T) < S_1^i$ then (T, S) is obtained by shifting cells of S to the left.

Recall also from Part I, 2.4.2 that given a tableau T we set $T^\dagger$ to be a transposed tableau. For example if T is the tableau above then

$$T^\dagger = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 5 & 6 & \\ \hline 4 & 6 & & \\ \hline 8 & & & \\ \hline 9 & & & \\ \hline \end{array}$$

2.3. We begin with the result $(**)$ from 1.7. This is a straightforward corollary of [2, 2.13]. Here we prove it for completeness.

Theorem. *For any $\mathcal{I} \subset \Pi$ one has*

$$\overline{\mathcal{V}}_{\mathcal{I}} = \bigcup_{\tau(\mathcal{V}) \supseteq \mathcal{I}} \mathcal{V}.$$

Proof.

First of all note that for any $w \in W$ one has $w \stackrel{D}{\geq} w_{\mathcal{I}}$ if and only if $\tau(w) \supseteq \mathcal{I}$. Thus

$$\overline{\mathcal{V}}_{\mathcal{I}} \supseteq \bigcup_{\tau(\mathcal{V}) \supseteq \mathcal{I}} \mathcal{V}.$$

Moreover since $\overline{\mathbf{B}(\mathbf{n} \cap {}^{w_{\mathcal{I}}}\mathbf{n})} = \mathbf{n} \cap {}^{w_{\mathcal{I}}}\mathbf{n}$ one has $\mathcal{V} \stackrel{G}{\geq} \mathcal{V}_{\mathcal{I}}$ if and only if $\mathcal{V} \stackrel{D}{\geq} \mathcal{V}_{\mathcal{I}}$. Hence $\mathcal{V} \stackrel{G}{\geq} \mathcal{V}_{\mathcal{I}}$ if and only if $\tau(\mathcal{V}) \supseteq \mathcal{I}$.

On the other hand

$$\overline{\mathcal{V}}_{\mathcal{I}} = \mathfrak{m}_{\mathcal{I}} = \coprod_{\mathcal{O} \geq \mathcal{O}_{\mathcal{I}}} \mathcal{O} \cap \mathfrak{m}_{\mathcal{I}}$$

So to complete the proof we must show that for any $\mathcal{O} > \mathcal{O}_{\mathcal{I}}$ any irreducible component of $\mathfrak{m}_{\mathcal{I}} \cap \mathcal{O}$ is an orbital variety. By [6, p. 456, last corollary] $\mathfrak{m}_{\mathcal{I}} \cap \mathcal{O}$ is equidimensional. By Part I, 4.1.8 it contains at least one orbital variety and so for every irreducible component $\mathcal{V}$ of $\mathcal{O} \cap \mathfrak{m}_{\mathcal{I}}$ one has $\dim \mathcal{V} = \frac{1}{2} \dim \mathcal{O}$. Yet $\mathcal{V} \subset \mathcal{O} \cap \mathbf{n}$, which also has dimension $\frac{1}{2} \dim \mathcal{O}$. Hence $\mathcal{V}$ is an orbital variety associated to $\mathcal{O}$. ■

2.4. Given a tableau T recall from 1.7 that $r_T(u)$ denotes the number of the row u belongs to.

Given $\mathcal{I} \subset \Pi$. Set $T_{\mathcal{I}} := T_{\mathcal{V}_{\mathcal{I}}}$. Since $T_{\mathcal{I}}$ is the minimal (both in the geometric and the Duflo orders) tableau such that $\tau(T) = \mathcal{I}$ one has

$$r_{T_{\mathcal{I}}}(i) = \begin{cases} 1, & \text{if } i = 1 \text{ or } \alpha_{i-1} \notin \mathcal{I}, \\ r_{T_{\mathcal{I}}}(i-1) + 1, & \text{if } \alpha_{i-1} \in \mathcal{I}. \end{cases}$$

A useful way to present $T_{\mathcal{I}}$ is provided in [2, 2.12]. Partition $\{1, 2, \dots, n\}$ into connected subsets $C_j := \{b_j, b_j + 1, \dots, b_{j+1} - 1\}$ by choosing a strictly increasing sequence $1 = b_1 < b_2 < \dots < b_{l+1} = n + 1$. Setting $\mathcal{I} = \{\alpha_i \mid i, i + 1 \text{ belong to some } C_j\}$ defines a bijection between the set of all such partitions and the set of subsets of Π . Given $\mathcal{I} \subset \Pi$, let $\{C_i^{\mathcal{I}} : i = 1, 2, \dots, l\}$ be the corresponding connected subsets which we view as columns. (Sometimes we may omit the $\mathcal{I}$ superscript.) Then in the notation of 2.2 we have $T_{\mathcal{I}} = (C_1^{\mathcal{I}}, C_2^{\mathcal{I}}, \dots, C_l^{\mathcal{I}})$. Of course this involves some sliding of boxes to the left. However there are some advantages in this presentation. For example $T_{\mathcal{I}} - 1$ is obtained by simply replacing $C_1^{\mathcal{I}}$ by $C_1^{\mathcal{I}} - 1$ and $T_{\mathcal{I}} - n$ is obtained by simply replacing $C_l^{\mathcal{I}}$ by $C_l^{\mathcal{I}} - n$.

We call $C_i^{\mathcal{I}}$ the i -th chain of $T_{\mathcal{I}}$ and $T_{\mathcal{I}} = (C_1^{\mathcal{I}}, C_2^{\mathcal{I}}, \dots, C_l^{\mathcal{I}})$ the chain form of $T_{\mathcal{I}}$.

Set $c_i := |C_i|$ and $\sigma_i := C_i^{c_i}$ which is the largest entry in C_i .

For example consider $\mathcal{I} = \{\alpha_1, \alpha_4, \alpha_5, \alpha_6, \alpha_9\}$ in $\mathfrak{sl}_{10}$. Then

$$T_{\mathcal{I}} = \begin{array}{|c|c|c|c|c|} \hline 1 & 3 & 4 & 8 & 9 \\ \hline 2 & 5 & 10 & & \\ \hline 6 & & & & \\ \hline 7 & & & & \\ \hline \end{array}$$

or in chain form $T_{\mathcal{I}} = (C_1, C_2, C_3, C_4, C_5)$ where $C_1 = \{1, 2\}$, $C_2 = \{3\}$, $C_3 = \{4, 5, 6, 7\}$, $C_4 = \{8\}$ and $C_5 = \{9, 10\}$. Here one has $c_1 = c_5 = 2$, $c_2 = c_4 = 1$ and $c_3 = 4$. As well $\sigma_1 = 2$, $\sigma_2 = 3$, $\sigma_3 = 7$, $\sigma_4 = 8$, $\sigma_5 = 10$.

2.5. Recall the notion $w_{\mathcal{I}}$ from 1.4. By 2.1 one has $\alpha_i \in \tau(w_{\mathcal{I}})$ if and only if $p_{w_{\mathcal{I}}}(i) > p_{w_{\mathcal{I}}}(i+1)$ and by 1.4 (*) we get $\alpha_i \in \tau(w_{\mathcal{I}})$ if and only if $\alpha_i \in \mathcal{I}$. Thus $p_{w_{\mathcal{I}}}(i) > p_{w_{\mathcal{I}}}(i+1)$ if and only if $\alpha_i \in \mathcal{I}$ which can be formulated as

Proposition ([2, 2.12]). *For all $\mathcal{I} \subset \Pi$, the word form of $w_{\mathcal{I}}$ is given by*

$$w_{\mathcal{I}} = [\overline{C_1^{\mathcal{I}}}, \overline{C_2^{\mathcal{I}}}, \dots, \overline{C_l^{\mathcal{I}}}].$$

2.6. For $s, t : 1 \leq s < t \leq l$ set $\hat{T}_{s,t} := (C_s, C_{s+1}, \dots, C_t)$.

For $i : 2 \leq i \leq l$ and for some $j : 1 \leq j \leq c_i$ set $C_i^{1,j-1} := (C_i^1, C_i^2, \dots, C_i^{j-1})$ and $C_i(j) := (C_i^j, \dots, C_i^{c_i})$. We call $C_i(j)$ a j -tail of the chain C_i . If there exist $s < i$ such

that $c_s \geq j$ then set

$$T' = \left(\begin{array}{c} (\hat{T}_{1,i-1})^{1,c_s}, C_i^{1,j-1} \\ (\hat{T}_{1,i-1})^{c_s+1,\infty}, C_i(j) \end{array} \right) \quad \text{and} \quad T_{\mathcal{I}}(c_s \swarrow C_i(j)) = (T', \hat{T}_{i+1,l}). \quad (*)$$

Note that T' is obtained from $\hat{T}_{1,i}$ just by moving j -tail of the last chain down to the part $(\hat{T}_{1,i-1})^{c_s+1,\infty}$. Since $|(\hat{T}_{1,i-1})^{c_s+1}| < |(\hat{T}_{1,i-1})^{c_s}|$ we get that T' and respectively $T_{\mathcal{I}}(c_s \swarrow C_i(j))$ are Young tableaux.

For example let $\mathcal{I} = \{\alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_7, \alpha_8\} \subset \Pi_{1,10}$ that is

$$T_{\mathcal{I}} = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 4 & 8 & \\ \hline 5 & 9 & & \\ \hline 6 & & & \\ \hline \end{array}$$

Then

$$T_{\mathcal{I}}(2 \swarrow C_2(2)) = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 8 & & \\ \hline 4 & 9 & & \\ \hline 5 & & & \\ \hline 6 & & & \\ \hline \end{array} \quad ; \quad T_{\mathcal{I}}(2 \swarrow C_3(2)) = \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 4 & & \\ \hline 5 & 8 & & \\ \hline 6 & 9 & & \\ \hline \end{array} \quad ;$$

$$T_{\mathcal{I}}(3 \swarrow C_4(1)) = \begin{array}{|c|c|c|} \hline 1 & 3 & 7 \\ \hline 2 & 4 & 8 \\ \hline 5 & 9 & \\ \hline 6 & 10 & \\ \hline \end{array}.$$

2.7. Consider $\hat{T}_{1,i-1}$, that is, look to the left of chain C_i . Set

$$\begin{aligned} \text{next}_l(i) &:= \begin{cases} \min\{c_j \mid j < i, c_j \geq c_i\}, & \text{if there exists } j < i \text{ such that } c_j \geq c_i; \\ 0, & \text{otherwise.} \end{cases} \\ \text{prev}_l(i) &:= \begin{cases} \max\{c_j \mid j < i, c_j \leq c_i\}, & \text{if there exists } j < i \text{ such that } c_j \leq c_i; \\ n, & \text{otherwise.} \end{cases} \\ \text{sprev}_l(i) &:= \begin{cases} \max\{c_j \mid j < i, c_j < c_i\}, & \text{if there exists } j < i \text{ such that } c_j < c_i; \\ n, & \text{otherwise.} \end{cases} \end{aligned}$$

Consider $\hat{T}_{i+1,l}$ that is look to the right of chain C_i . Set

$$\begin{aligned} \text{snext}_r(i) &:= \begin{cases} \min\{c_j \mid j > i, c_j > c_i\}, & \text{if there exists } j > i \text{ such that } c_j > c_i; \\ n, & \text{otherwise.} \end{cases} \\ \text{sprev}_r(i) &:= \begin{cases} \max\{c_j \mid j > i, c_j < c_i\}, & \text{if there exists } j > i \text{ such that } c_j < c_i; \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

We use the nomenclature “sprev” (resp. “snext”) to emphasize that this is a strictly previous (resp. a strictly next) number.

Set

$$T_{\mathcal{I}}(i, \text{next}) := \begin{cases} T_{\mathcal{I}}(\text{next}_l(i) \swarrow C_i(c_i)), & \text{if } \text{next}_l(i) \neq 0, \\ \emptyset, & \text{otherwise.} \end{cases}$$

That is consider $\hat{T}_{1,i}$ and move the box with σ_i , which is the largest number of $\hat{T}_{1,i}$, down to the first possible row (if such row exists), then add to the new tableau $\hat{T}_{i+1,l}$.

Set

$$T_{\mathcal{I}}(i, \text{prev}) = \begin{cases} T_{\mathcal{I}}(\text{sprev}_l(i) \swarrow C_i(\text{sprev}_l(i))), & \text{if } \text{prev}_l(i) = \text{sprev}_l(i) \neq n, \\ \emptyset, & \text{otherwise.} \end{cases}$$

That is consider $\hat{T}_{1,i}$ and move the smallest possible tail of the last column one row down (if this is possible), then add to the new tableau $\hat{T}_{i+1,l}$.

Set $T_{\mathcal{I}}(i) := \{T_{\mathcal{I}}(i, \text{next}), T_{\mathcal{I}}(i, \text{prev})\}$.

For example consider $T_{\mathcal{I}}$ from 2.6. Then

$$\begin{aligned} T_{\mathcal{I}}(2) &= \left\{ \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 8 & & \\ \hline 4 & 9 & & \\ \hline 5 & & & \\ \hline 6 & & & \\ \hline \end{array} \right\}, & T_{\mathcal{I}}(3) &= \left\{ \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 4 & 8 & \\ \hline 5 & & & \\ \hline 6 & & & \\ \hline 9 & & & \\ \hline \end{array} \right\}, & \left\{ \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 4 & & \\ \hline 5 & 8 & & \\ \hline 6 & 9 & & \\ \hline \end{array} \right\}, \\ \\ T_{\mathcal{I}}(4) &= \left\{ \begin{array}{|c|c|c|} \hline 1 & 3 & 7 \\ \hline 2 & 4 & 8 \\ \hline 5 & 9 & 10 \\ \hline 6 & & \\ \hline \end{array} \right\}. \end{aligned}$$

Remark 1. For $i > 1$ one has $T_{\mathcal{I}}(i) \neq \emptyset$.

Remark 2. Define

$$\hat{T}_{\mathcal{I}}(i, \text{prev}) := \begin{cases} T_{\mathcal{I}}(i, \text{prev}_l(i), \text{prev}_l(i)), & \text{if } \text{prev}_l(i) \neq n, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Then

$$\hat{T}_{\mathcal{I}}(i, \text{prev}) := \begin{cases} T_{\mathcal{I}}(i, \text{next}), & \text{if } \text{prev}_l(i) = c_i, \\ T_{\mathcal{I}}(i, \text{prev}), & \text{otherwise.} \end{cases}$$

That is why we define $T_{\mathcal{I}}(i, \text{prev})$ with the help of $\text{sprev}_l(i)$ and not of $\text{prev}_l(i)$.

2.8. Let $\mathcal{D}_G(T)$ denote the set of geometric descendants of T . Now we can formulate the main theorem of this paper.

Theorem. For $T_{\mathcal{I}}$ the set of descendants is defined by

$$\mathcal{D}_G(T_{\mathcal{I}}) = \{T_{\mathcal{I}}(i, \text{next}) \mid \text{next}_l(i) = c_i \text{ or } \langle \text{next}_l(i) > c_i \text{ and } \text{snext}_r(i) > \text{next}_l(i) \rangle\} \cup \{T_{\mathcal{I}}(i, \text{prev}) \mid \text{sprev}_l(i) < n \text{ and } \text{sprev}_r(i) < \text{sprev}_l(i)\}.$$

In our example from 2.6 one has

$$\mathcal{D}_G(T_{\mathcal{I}}) = \left\{ \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 4 & 8 & \\ \hline 5 & & & \\ \hline 6 & & & \\ \hline 9 & & & \\ \hline \end{array} \right\}, \quad \left\{ \begin{array}{|c|c|c|c|} \hline 1 & 3 & 7 & 10 \\ \hline 2 & 4 & & \\ \hline 5 & 8 & & \\ \hline 6 & 9 & & \\ \hline \end{array} \right\}, \quad \left\{ \begin{array}{|c|c|c|} \hline 1 & 3 & 7 \\ \hline 2 & 4 & 8 \\ \hline 5 & 9 & 10 \\ \hline 6 & & \\ \hline \end{array} \right\}.$$

Consider $i : 2 \leq i \leq l$. Note that for any $T \in T_{\mathcal{I}}(i)$ one has $\tau(T) \supset \mathcal{I}$ thus $T \stackrel{G}{>} T_{\mathcal{I}}$.

The proof of the theorem consists of the following steps:

(i) We show that

$$\mathcal{D}_G(T_{\mathcal{I}}) \subset \bigcup_{i=2}^l T_{\mathcal{I}}(i).$$

(ii) We show that $T \in T_{\mathcal{I}}(l)$ is always in $\mathcal{D}_G(T_{\mathcal{I}})$.

(iii) Finally we consider $T \in T_{\mathcal{I}}(i)$ for $i : 1 < i < l$. Note that if $\text{next}_l(i) = c_i$ then $T_{\mathcal{I}}(i) = \{T_{\mathcal{I}}(i, \text{next})\}$ and $\text{codim}_{\mathfrak{m}_{\mathcal{I}}} \mathcal{V}_{T_{\mathcal{I}}(i, \text{next})} = 1$, thus $T_{\mathcal{I}}(i, \text{next}) \in \mathcal{D}_G(T_{\mathcal{I}})$ just by the dimension consideration. We must consider only $T \in T_{\mathcal{I}}(i)$ such that $\text{codim}_{\mathfrak{m}_{\mathcal{I}}} \mathcal{V}_T > 1$, that is $T_{\mathcal{I}}(i, \text{next})$ when $\text{next}_l(i) > c_i$ and $T_{\mathcal{I}}(i, \text{prev})$ if it exists. To deal with these cases we use order preserving involution ψ on $\mathbf{T}_n$ induced by the non-trivial involution of the Dynkin diagram of $\mathfrak{sl}_n$. With the help of ψ we first show that in case $\text{next}_l(i) > c_i$ one has $T_{\mathcal{I}}(i, \text{next}) \in \mathcal{D}_G(T_{\mathcal{I}})$ iff $\text{snext}_r(i) > \text{next}_l(i)$. Then we use ψ again to show that $T_{\mathcal{I}}(i, \text{prev}) \in \mathcal{D}_G(T_{\mathcal{I}})$ iff $\text{sprev}_r(i) < \text{sprev}_l(i)$.

3. Proof of the theorem on descendants

3.1. For $1 \leq i < j \leq n$ set $\langle i, j \rangle := \{k\}_{k=i}^j$. Recall the notion of projection $\pi_{i,j} : \mathbf{T}_n \rightarrow \mathbf{T}_{\langle i,j \rangle}$ obtained through jeu de taquin applied to entries of $T \in \mathbf{T}_n$ not lying in $\langle i, j \rangle$ as it is defined in I, 2.4.16.

3.2. Let us consider the projections $\pi_{i,j}$ of $T \in T_{\mathcal{I}}(i)$. As noted in 2.4 the tableau $\pi_{1,n-1}(T_{\mathcal{I}})$ is obtained from $T_{\mathcal{I}}$ by eliminating n from C_l . It corresponds to the subset $\mathcal{I}_n := \mathcal{I} \cap \Pi_{1,n-1}$ with $C_i^{\mathcal{I}_n} = C_i^{\mathcal{I}} : i < l$ and $C_l^{\mathcal{I}_n} = C_l^{\mathcal{I}} - n$. Similarly $\pi_{2,n}(T_{\mathcal{I}})$ is obtained from $T_{\mathcal{I}}$ by eliminating 1 from C_1 and corresponds to the subset $\mathcal{I}_1 := \mathcal{I} \cap \Pi_{2,n}$ with $C_i^{\mathcal{I}_1} = C_i^{\mathcal{I}} : i > 1$ and $C_1^{\mathcal{I}_1} = C_1^{\mathcal{I}} - 1$.

If $T_{\mathcal{I}}(i, \text{next}) \neq \emptyset$ set $p(i, \text{next}) := \max\{s \mid s < i, c_s = \text{next}_l(i)\}$. If $T_{\mathcal{I}}(i, \text{prev}) \neq \emptyset$ set $p(i, \text{prev}) := \max\{s \mid s < i, c_s = \text{sprev}_l(i)\}$.

Lemma. For $i : i > 1$ consider $T \in T_{\mathcal{I}}(i)$ and fix

$$p(T) = \begin{cases} p(i, \text{next}), & \text{if } T = T_{\mathcal{I}}(i, \text{next}) \\ p(i, \text{prev}), & \text{otherwise} \end{cases}$$

Then

- (i) $\pi_{1,n-1}(T) \in T_{\mathcal{I}_n}(i)$, if $i < l$.
- (ii) If $T_{\mathcal{I}}(l, \text{next}) \neq \emptyset$ then $\pi_{1,n-1}(T_{\mathcal{I}}(l, \text{next})) = T_{\mathcal{I}_n}$.
- (iii) $\pi_{1,n-1}(T_{\mathcal{I}}(l, \text{prev})) \in T_{\mathcal{I}_n}(l)$.
- (iv) $\pi_{2,n}(T) \in T_{\mathcal{I}_1}(i)$, if $p(T) > 1$ or $p(T) = 1$ and $c_1 > c_i$.
- (v) $\pi_{2,n}(T) = T_{\mathcal{I}_1}$, if $p(T) = 1$ and $c_1 \leq c_i$.

Proof.

(i) is an immediate consequence of $\sigma_i < n$, for $i < l$.

(ii) is an immediate consequence of the definition of $T_{\mathcal{I}}(l, \text{next})$ (which is defined by moving the box containing n down) and of $\pi_{1,n-1}(T)$ (which is defined by eliminating box containing n from T).

(iii) is an immediate consequence of $C_l^{\text{sprev}(l)} < n$.

To prove (iv) note that $T_{\mathcal{I}_1}$ is obtained by replacing C_1 by $C_1 - 1$. If $p(T) > 1$ then the operation of moving the tail commutes with this replacing. If $p(T) = 1$ and $c_1 > c_i$ then $T = T_{\mathcal{I}}(i, \text{next})$ and σ_i goes from c_i -th row to $(c_1 + 1)$ -th row in forming $T_{\mathcal{I}}(i, \text{next})$, but then goes to c_1 -th row in forming $\pi_{2,n}(T_{\mathcal{I}}(i, \text{next}))$. However if $c_1 > c_i$ this is exactly what happens in forming $T_{\mathcal{I}_1}(i, \text{next})$ from $T_{\mathcal{I}_1}$. Hence (iv).

Finally if $c_i > c_1 > c_j$, $\forall j : 1 < j < i$, that is under hypothesis of (v) for $T_{\mathcal{I}}(i, \text{prev})$, one has $T_2^{c_1} = C_i^{c_1}$ and it is pushed back to c_1 -th row in forming $\pi_{2,n}(T)$ which is hence $T_{\mathcal{I}_1}$. ■

3.3. Recall notation $p_w(i)$ from 2.1. In what follows we need the following simple

Lemma. *Given $x, z \in \mathbf{S}_n$ such that*

- (i) $\pi_{1,s}(x) = \pi_{1,s}(z)$
- (ii) $p_x(n) < p_x(n-1) < \cdots < p_x(s+1), p_z(n) < p_z(n-1) < \cdots < p_z(s+1)$
- (iii) $p_x(j) \leq p_z(j) \quad \forall j : s+1 \leq j \leq n.$

Then $\mathbf{n} \cap^x \mathbf{n} \subset \mathbf{n} \cap^z \mathbf{n}.$

Proof.

Recall from 2.1 that $\alpha_{i,j} \in S(w)$ if and only if $p_w(i) > p_w(j)$. Applying this proposition to all 3 hypotheses we get

- (i) implies that for any $i, j : i < j \leq s$ one has $\alpha_{i,j} \in S(x)$ iff $\alpha_{i,j} \in S(z)$;
- (ii) implies that for any $i, j : s+1 \leq i < j$ one has $\alpha_{i,j} \in S(x)$ and $\alpha_{i,j} \in S(z)$;
- (iii) implies that for any $i, j : i \leq s, j \geq s+1$ one has $\alpha_{i,j} \in S(z)$ implies $\alpha_{i,j} \in S(x)$.

Thus $S(z) \subset S(x)$ which is equivalent to $\mathbf{n} \cap^x \mathbf{n} \subset \mathbf{n} \cap^z \mathbf{n}.$ ■

3.4. Given tableaux P, Q such that $\langle P \rangle \cap \langle Q \rangle = \emptyset$, recall the notation (P, Q) from 2.2 or Part I, 2.4.3 and set $\binom{P}{Q} := (P, Q)^\dagger$.

In what follows we need the lemma which is a kind of variation of Part I, 3.2.3 (v),(vi).

Lemma. *Given tableaux P and Q such that for any $p \in \langle P \rangle$ and any $q \in \langle Q \rangle$ one has $p < q$. For any $x \in \mathcal{C}_P$ and $y \in \mathcal{C}_Q$ one has*

- (i) *Set $w = [x, y]$. Then $T(w) = (P, Q)$.*
- (ii) *Set $w = [y, x]$. Then $T(w) = \binom{P}{Q}$.*

Proof.

To show (i) it is enough to show that for any row i of P , for any $p \in P^i$ one has $r_{T(w)}(p) = i$ and for any row i of Q , for any $q \in Q^i$ one has $r_{T(w)}(q) = i$. Let $x = [x_1, \dots, x_k]$ and $y = [y_1, \dots, y_m]$.

Recall notation of RS procedure from Part I, 2.4.6. One has ${}_k T(w) = T(x) = P$ so that $r_{{}_k T(w)}(p) = i$. For any y_j one has $y_j > p$ thus RS insertion does not knock down p so that $r_{T(w)}(p) = i$. On the other hand again since any y_j does not knock down any $x_{j'}$ one has that it knocks down exactly the same Q_j^i as in $T(y)$. Thus if $q \in Q^i$ then $r_{T(w)}(q) = i$.

To show (ii) we use Schensted-Schützenberger theorem (cf. Part I, 2.4.15) claiming $T(\overline{w}) = (T(w))^\dagger$. First of all this theorem together with part (i) gives us that $T([\overline{x}, \overline{y}]) =$

$(P^\dagger, Q^\dagger)$. Applying the theorem again we get

$$T([y, x]) = T([\overline{x}, \overline{y}]) = (P^\dagger, Q^\dagger)^\dagger = \begin{pmatrix} P \\ Q \end{pmatrix}.$$

■

3.5. Now we are ready to show that $\mathcal{D}_G(T) \subset \{T_{\mathcal{I}}(i)\}_{i=2}^l$. Let $T \stackrel{G}{>} T_{\mathcal{I}}$ and let m be minimal integer such that $r_T(m) > r_{T_{\mathcal{I}}}(m)$. Let i be the number of the chain m belongs to, that is $m \in < C_i^{\mathcal{I}} >$. If $T_{\mathcal{I}}(i)$ contains only one element let $T_{\mathcal{I}}[m]$ denote this element. If $T_{\mathcal{I}}(i) = \{T_{\mathcal{I}}(i, \text{next}), T_{\mathcal{I}}(i, \text{prev})\}$ then

$$T_{\mathcal{I}}[m] = \begin{cases} T_{\mathcal{I}}(i, \text{prev}), & \text{if } m \leq C_i^{\text{sprev}_l(i)} \\ T_{\mathcal{I}}(i, \text{next}), & \text{otherwise} \end{cases}$$

Proposition. *Let $\overline{\mathcal{V}}_T \subsetneq \mathfrak{m}_{\mathcal{I}}$ and let m be minimal integer such that $r_T(m) > r_{T_{\mathcal{I}}}(m)$. Then $\overline{\mathcal{V}}_T \subset \overline{\mathcal{V}}_{T_{\mathcal{I}}[m]}$. In particular $\mathcal{D}_G(T) \subset \{T_{\mathcal{I}}(i)\}_{i=2}^l$.*

Proof.

This is trivially true for $\mathfrak{sl}_3$. Assume this is true for $n-1$ and show this for n .

Assume $i < l$. Consider $\pi_{1,n-1}(T_{\mathcal{I}}[m])$. By 3.2 (i) in that case $\pi_{i,n-1}(T_{\mathcal{I}}[m]) = T_{\mathcal{I}_n}[m]$. By Part I, 4.1.2 one has $\pi_{1,n-1}(\overline{\mathcal{V}}_T) \subset \mathfrak{m}_{\mathcal{I}_n}$. As well, since $m < n$ in that case, one has $r_{\pi_{1,n-1}(T)}(m) = r_T(m) > r_{T_{\mathcal{I}_n}}(m)$. Thus by induction assumption $\overline{\mathcal{V}}_{\pi_{1,n-1}(T)} \subset \overline{\mathcal{V}}_{T_{\mathcal{I}_n}[m]}$.

On the other hand we can write $\mathfrak{m}_{\mathcal{I}}$ as a Cartesian product:

$$\mathfrak{m}_{\mathcal{I}} = \mathfrak{m}_{\mathcal{I}_n} \times \sum_{j=1}^{\sigma_{l-1}} X_{\alpha_{j,n}}.$$

Thus for any $\overline{\mathcal{V}}_T \subset \mathfrak{m}_{\mathcal{I}}$ one has

$$\overline{\mathcal{V}}_T \subset \overline{\mathcal{V}}_{\pi_{1,n-1}(T)} \times \sum_{j=1}^{\sigma_{l-1}} X_{\alpha_{j,n}}.$$

For any $i : 1 < i < l$ consider $w = w_r(\pi_{1,\sigma_{l-1}}(T_{\mathcal{I}}[m]))$ and $x = \overline{C}_l$. By 3.4 (i) one has $Q([w, x]) = T_{\mathcal{I}}[m]$.

By Part I, 2.2.4 one has $\alpha_{j,n} \in S([w, x])$ if and only if $j \in C_l$. Thus $\overline{\mathcal{V}}_{T_{\mathcal{I}}[m]} = \overline{\mathcal{V}}_{T_{\mathcal{I}_n}[m]} \times \sum_{j=1}^{\sigma_{l-1}} X_{\alpha_{j,n}}$. Hence if $i < l$ then

$$\overline{\mathcal{V}}_T \subset \overline{\mathcal{V}}_{\pi_{1,n-1}(T)} \times \sum_{j=1}^{\sigma_{l-1}} X_{\alpha_{j,n}} \subset \overline{\mathcal{V}}_{T_{\mathcal{I}_n}[m]} \times \sum_{j=1}^{\sigma_{l-1}} X_{\alpha_{j,n}} = \overline{\mathcal{V}}_{T_{\mathcal{I}}[m]}$$

If $i = l$ let us show that there exists $y : T(y) = T$ and $z : T(z) = T_{\mathcal{I}}[m]$ such that $\mathfrak{n} \cap^y \mathfrak{n} \subset \mathfrak{n} \cap^z \mathfrak{n}$. Indeed let $y = w_r(T)$ and $z = w_r(T_{\mathcal{I}}[m])$. Note that

- (i) $\pi_{1,m-1}(T) = \pi_{1,m-1}(T_{\mathcal{I}}[m])$ therefore $\pi_{1,m-1}(y) = \pi_{1,m-1}(z)$.
- (ii) $r_T(n) > r_T(n-1) > \dots > r_T(m)$ and $r_{T_{\mathcal{I}}[m]}(n) > r_{T_{\mathcal{I}}[m]}(n-1) > \dots > r_{T_{\mathcal{I}}[m]}(m)$ therefore $p_y(n) < p_y(n-1) < \dots < p_y(m)$ and $p_z(n) < p_z(n-1) < \dots < p_z(m)$.
- (iii) (a) If $m \leq C_l^{\text{sprev}_l(l)}$ then for any $s \geq m$ one has

$$r_{T_{\mathcal{I}}[m]}(s) = \begin{cases} r_{T_{\mathcal{I}}}(s), & \text{if } s < C_l^{\text{sprev}_l(l)}, \\ r_{T_{\mathcal{I}}}(s) + 1, & \text{otherwise.} \end{cases}$$

On the other hand for any $s \geq m$ one has $r_T(s) \geq r_{T_{\mathcal{I}}}(s) + 1$ just by $\{\alpha_s\}_{s=m}^{n-1} \subset \tau(T_{\mathcal{I}}) \subset \tau(T)$ and by the condition $r_T(m) \geq r_{T_{\mathcal{I}}}(m) + 1$. Thus $p_y(s) \leq p_z(s)$.

- (b) If $m > C_l^{\text{sprev}_l(l)}$ then

$$r_{T_{\mathcal{I}}[m]}(s) = \begin{cases} r_{T_{\mathcal{I}}}(s), & \text{if } s < n, \\ \text{snext}_l(l) + 1, & \text{if } s = n. \end{cases}$$

On the other hand since $|T_{\mathcal{I}}^{\text{sprev}_l(l)+1}| = \dots = |T_{\mathcal{I}}^{c_l}|$ and $|T_{\mathcal{I}}^{c_l+1}| = \dots = |T_{\mathcal{I}}^{\text{snext}_l(l)}| = |T_{\mathcal{I}}^{c_l}| - 1$ one has that $r_T(m) \geq \text{snext}_l(l) + 1$ and for any $s \geq m$ one has $r_T(s) \geq \text{snext}_l(l) + 1 + (s - m)$ exactly by the same reasoning as in (a). Thus again $p_y(s) \leq p_z(s)$.

In both cases we get the hypothesis (iii) of 3.3. Therefore by 3.3 we get the result. ■

Remark. Recall that $T_{\mathcal{I}}[m] \in T_{\mathcal{I}}(i)$. Let $w = w_r(\pi_{1,\sigma_i}(T_{\mathcal{I}}[m]))$ and

$$x = \begin{cases} [\overline{C}_{i+1}, \dots, \overline{C}_l] & \text{if } i < l, \\ \emptyset, & \text{if } i = l. \end{cases}$$

then by 3.4 (i) one has $Q[w, x] = T_{\mathcal{I}}[m]$ and the same proof shows that

$$\mathbf{n} \cap^{w_r(T)} \mathbf{n} \subset \mathbf{n} \cap^{[w, x]} \mathbf{n},$$

that is $T \stackrel{\text{D}}{>} T_{\mathcal{I}}[m]$. In particular this means that $T \stackrel{\text{G}}{>} T_{\mathcal{I}}[m]$ if and only if $T \stackrel{\text{D}}{>} T_{\mathcal{I}}[m]$. The interesting question is whether this is true for any $T \in \{T_{\mathcal{I}}(i)\}_{i=2}^l$, that is whether for any $T \in \{T_{\mathcal{I}}(i)\}_{i=2}^l$ one has $S \stackrel{\text{G}}{>} T$ if and only if $S \stackrel{\text{D}}{>} T$.

3.6. We have completed step (i) of the proof. Now we prove step (ii), namely

Proposition. *Given $T_{\mathcal{I}} = (C_1, \dots, C_l)$.*

- (i) *If $T_{\mathcal{I}}(l, \text{next}) \neq \emptyset$ then $T_{\mathcal{I}}(l, \text{next}) \in \mathcal{D}(T_{\mathcal{I}})$.*
- (ii) *If $T_{\mathcal{I}}(l, \text{prev}) \neq \emptyset$ then $T_{\mathcal{I}}(l, \text{prev}) \in \mathcal{D}(T_{\mathcal{I}})$.*

Proof.

Set $\text{sh}(T_{\mathcal{I}}) = \lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$.

(i) Assume that $T_{\mathcal{I}}(l, \text{next}) \neq \emptyset$ and put $T := T_{\mathcal{I}}(l, \text{next})$, $l' := p(l, \text{next})$ and $\text{sh}(T) := \mu = (\mu_1, \dots, \mu_k)$. (If the number of rows in T is greater by 1 than the number of rows in $T_{\mathcal{I}}$ we suppose that indeed $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{k-1})$ and add $\lambda_k = 0$ to λ to get the same length.) One has

$$\mu_j = \begin{cases} \lambda_j, & \text{if } j \neq c_l, c_{l'} + 1, \\ \lambda_{c_l} - 1, & \text{if } j = c_l, \\ \lambda_{c_{l'}+1} + 1, & \text{if } j = c_{l'} + 1. \end{cases}$$

If $c_{l'} = c_l$ then $\mathcal{O}_\mu$ is a descendant of $\mathcal{O}_\lambda$ by Part I, 2.3.3. Thus $\mathcal{V}_T$ is a descendant of $\mathcal{V}_{\mathcal{I}}$. If for any $j \neq l'$ one has $c_j \neq c_{l'}$ then $\lambda_{c_l+1} = \dots = \lambda_{c_{l'}} = \lambda_{c_l} - 1$ and $\lambda_{c_{l'}+1} = \lambda_{c_l} - 2$ so that again $\mathcal{O}_\mu$ is a descendant of $\mathcal{O}_\lambda$ by Part I, 2.3.3. Thus again $\mathcal{V}_T$ is a descendant of $\mathcal{V}_{\mathcal{I}}$.

Assume that $c_{l'} > c_l$ and there exist $j \neq l'$ such that $c_j = c_{l'}$. Then there exist the unique orbit $\mathcal{O}_\nu$ such that $\overline{\mathcal{O}}_\mu \subsetneq \overline{\mathcal{O}}_\nu \subsetneq \overline{\mathcal{O}}_\lambda$ where $\nu = (\nu_1, \dots, \nu_k)$ is defined by

$$\nu_j = \begin{cases} \lambda_j, & \text{if } j \neq c_{l'}, c_{l'} + 1, \\ \lambda_{c_{l'}} - 1, & \text{if } j = l', \\ \lambda_{c_{l'}+1} + 1, & \text{if } j = c_{l'} + 1. \end{cases}$$

Assume that there exists T' such that $\overline{\mathcal{V}}_T \subsetneq \overline{\mathcal{V}}_{T'} \subsetneq \overline{\mathcal{V}}_{\mathcal{I}}$. Then $\text{sh}(T') = \nu$. Consider $\pi_{1,n-1}$ of the three orbital variety closures. We get $\overline{\mathcal{V}}_{\pi_{1,n-1}(T)} \subset \overline{\mathcal{V}}_{\pi_{1,n-1}(T')} \subset \overline{\mathcal{V}}_{\pi_{1,n-1}(\mathcal{I})}$. Since $\pi_{1,n-1}(T) = \pi_{1,n-1}(T_{\mathcal{I}})$ one has $\pi_{1,n-1}(T') = \pi_{1,n-1}(T_{\mathcal{I}})$ which is impossible since $|(\pi_{1,n-1}(T_{\mathcal{I}}))^{c_{l'}}| = \lambda_{c_{l'}}$ and $|(\pi_{1,n-1}(T'))^{c_{l'}}| \leq \nu_{c_{l'}} = \lambda_{c_{l'}} - 1$.

(ii) The proof of (ii) is very similar to the proof of (i). Assume that $T_{\mathcal{I}}(l, \text{prev}) \neq \emptyset$ and put $T = T_{\mathcal{I}}(l, \text{prev})$, $l' := p(l, \text{prev})$ and $\text{sh}(T) := \mu = (\mu_1, \dots, \mu_k)$. (Again if the number of rows in T greater by 1 than the number of rows in $T_{\mathcal{I}}$ we again add $\lambda_k = 0$ to λ to get the same length.) One has

$$\mu_j = \begin{cases} \lambda_j, & \text{if } j \neq c_{l'}, c_l + 1, \\ \lambda_{c_{l'}} - 1, & \text{if } j = c_{l'}, \\ \lambda_{c_l+1} + 1, & \text{if } j = c_l + 1. \end{cases}$$

Note that by definition of $T_{\mathcal{I}}(l, \text{prev})$ we obtain

$$\lambda_{c_{l'}+1} = \dots = \lambda_{c_l} \quad \text{and} \quad \lambda_{c_l+1} = \lambda_{c_l} - 1. \quad (*)$$

The claim is trivially true for $n = 3$ so assume that in $\mathbf{T}_{n-1}$ one has that $T_{\mathcal{I}}(l, \text{prev})$ is a descendant of $T_{\mathcal{I}}$.

Again, if for any $j \neq l'$ one has $c_j \neq c_{l'}$ then $\mathcal{O}_\mu$ is a descendant of $\mathcal{O}_\lambda$ by Part I, 2.3.3, thus $\mathcal{V}_T$ is a descendant of $\mathcal{V}_{\mathcal{I}}$.

Again, assume that $c_{l'} < c_l$ and there exist $j \neq l'$ such that $c_j = c_{l'}$. Note that in that case in particular

$$\alpha_{n-1} \in \mathcal{I} \quad (**).$$

By (*) in that case there exist the unique orbit $\mathcal{O}_\nu$ such that $\overline{\mathcal{O}}_\mu \subsetneq \overline{\mathcal{O}}_\nu \subsetneq \overline{\mathcal{O}}_\lambda$ where $\nu = (\nu_1, \dots, \nu_k)$ is defined by

$$\nu_j = \begin{cases} \lambda_j, & \text{if } j \neq c_{l'}, c_{l'} + 1, \\ \lambda_{c_{l'}} - 1, & \text{if } j = l', \\ \lambda_{c_{l'}+1} + 1, & \text{if } j = c_{l'} + 1. \end{cases}$$

Assume that there exist T' such that $\overline{\mathcal{V}}_T \subsetneq \overline{\mathcal{V}}_{T'} \subsetneq \overline{\mathcal{V}}_{\mathcal{I}}$ then $\text{sh}(T') = \nu$. Consider $\pi_{1,n-1}$ of the three orbital variety closures. We get $\overline{\mathcal{V}}_{\pi_{1,n-1}(T)} \subset \overline{\mathcal{V}}_{\pi_{1,n-1}(T')} \subset \overline{\mathcal{V}}_{\mathcal{I}_n}$. By the induction assumption $\pi_{1,n-1}(T') = \pi_{1,n-1}(T)$ or $\pi_{1,n-1}(T') = T_{\mathcal{I}_n}$. The second situation is impossible since $|T_{\mathcal{I}_n}^{c_{l'}}| = \lambda_{c_{l'}}$ and $|(\pi_{1,n-1}(T'))^{c_{l'}}| \leq \nu_{c_{l'}} = \lambda_{c_{l'}} - 1$. Thus $\pi_{1,n-1}(T') = \pi_{1,n-1}(T)$. Set $\mu' = \text{sh}(\pi_{1,n-1}(T)) = \text{sh}(\pi_{1,n-1}(T'))$. Then

$$\mu'_j = \begin{cases} \lambda_j, & \text{if } j \neq c_{l'}, \\ \lambda_{c_{l'}} - 1, & \text{if } j = c_{l'}. \end{cases} = \begin{cases} \nu_j, & \text{if } j \neq c_{l'} + 1, \\ \nu_{c_{l'}+1} - 1, & \text{if } j = c_{l'} + 1. \end{cases}$$

Thus $r_{T'}(n) = c_{l'} + 1 \leq r_T(n-1) = c_l$. Since $\pi_{1,n-1}(T') = \pi_{1,n-1}(T)$ we get that $r_T(n-1) = r_{T'}(n-1)$ so that $\alpha_{n-1} \notin \tau(T')$ which contradicts to (**). ■

3.7. Note that 3.6 (i) can be easily generalized to any Young tableau T

Proposition. *Given $T \in \mathbf{T}_n$. Let $\text{sh}(T) = \lambda = (\lambda_1, \dots, \lambda_k)$. Let $i := r_T(n)$ and let $j = \min\{m \mid \lambda_m < \lambda_i - 1\}$. Then S obtained from T by moving box with n from row i to row j , or formally*

$$S := \begin{pmatrix} T^{1,i-1} \\ T^i - n \\ T^{i+1,j-1} \\ T^j + n \\ T^{j+1,\infty} \end{pmatrix},$$

is a geometric descendant of T .

The proof is exactly the same as the proof of 3.6 (i). In particular 3.6 and 3.7 show that $\mathcal{V}_S$ being a geometric descendant of $\mathcal{V}_T$ does not necessarily imply that $\mathcal{O}_S$ is a descendant of $\mathcal{O}_T$.

3.8. As it is formulated in theorem 2.8, it may happen that for a given $I : i < l$, $T_{\mathcal{I}}(i, \text{next})$ or $T_{\mathcal{I}}(i, \text{prev})$ is *not* a geometric descendant of $T_{\mathcal{I}}$. This is a difficult point. To understand it we need the non-trivial involution ψ of the Dynkin diagram of $\mathfrak{sl}_n$ defined by $\psi(\alpha_i) = \alpha_{n-i}$. This involution induces the involution ψ of $\mathfrak{sl}_n$ obtained by $\psi(X_{\alpha_{i,j}}) = X_{\alpha_{n+1-j, n+1-i}}$ and of its Weyl group $W = \mathbf{S}_n$ defined by $\psi(s_{\alpha_i}) = s_{\alpha_{n-i}}$. One has $\psi(w)(\psi(\alpha_{i,j})) = \psi(w(\alpha_{i,j}))$. The involution has a nice description on word presentations. Set $\psi_n : \{i\}_{i=1}^n \rightarrow \{i\}_{i=1}^n$ by $\psi_n(i) := n + 1 - i$.

Lemma. Let $w = [a_1, a_2, \dots, a_n]$ be a word presentation.

Then $\psi(w) = [\psi_n(a_n), \psi_n(a_{n-1}), \dots, \psi_n(a_1)]$.

Proof.

First of all let us show the assertion for $s_i = [1, 2, \dots, i-1, i+1, i, i+2, \dots, n]$

$$\begin{aligned} \psi(s_i) = s_{n-i} &= [1, 2, \dots, n-i-1, n-i+1, n-i, \dots, n] \\ &= [\psi_n(n), \dots, \psi_n(i+2), \psi_n(i), \psi_n(i+1), \psi_n(i-1), \dots, \psi_n(1)]. \end{aligned}$$

Recall from Part I, 1.9 the notion of $\ell(w)$ – that is the minimal length of w as an element of Weyl group written as a product of $s_\alpha : \alpha \in \Pi$. Assume that the assertion is true for any w of length $\ell(w) \leq p-1$ and show this for $w = s_{i_1} \cdots s_{i_p}$. Let $w' = s_{i_1} \cdots s_{i_{p-1}} = [a_1, \dots, a_n]$ in word presentation. Then $w = w' s_{i_p} = [b_1, \dots, b_n]$ where

$$b_j = \begin{cases} a_j, & \text{if } j \neq i_p, i_p + 1, \\ a_{i_p+1}, & \text{if } j = i_p, \\ a_{i_p}, & \text{if } j = i_p + 1. \end{cases}$$

One has

$$\begin{aligned} \psi(w) &= \psi(w') s_{n-i_p} \\ &= [\psi_n(a_n), \dots, \psi_n(a_{i_p}), \psi_n(a_{i_p+1}), \dots, \psi_n(a_1)] s_{n-i_p} \quad \text{by ind. assumption} \\ &= [\psi_n(a_n), \psi_n(a_{n-1}), \dots, \psi_n(a_{i_p+1}), \psi_n(a_{i_p}), \dots, \psi_n(a_1)] \\ &= [\psi_n(b_n), \psi_n(b_{n-1}), \dots, \psi_n(b_{i_p}), \psi_n(b_{i_p+1}), \dots, \psi_n(b_1)] \end{aligned}$$

■

In what follows we use the following notation. For $w = [a_1, \dots, a_m]$ such that $\langle w \rangle \subset \{i\}_{i=1}^n$ put $\psi(w) := [\psi_n(a_m), \psi_n(a_{m-1}), \dots, \psi_n(a_1)]$.

3.9. The involution ψ on W induces the (geometric and Duflo) order preserving involution on $\mathbf{T}_n$ obtained by $\psi(T(w)) := T(\psi(w))$. Indeed since this is an involution on $\mathfrak{sl}_n$ and on $\mathbf{B}$ and $\psi(\mathbf{n} \cap^w \mathbf{n}) = \mathbf{n} \cap^{\psi(w)} \mathbf{n}$ we obtain

$$\mathbf{n} \cap^w \mathbf{n} \subseteq \overline{\mathbf{B}(\mathbf{n} \cap^y \mathbf{n})} \quad \text{iff} \quad \mathbf{n} \cap^{\psi(w)} \mathbf{n} \subseteq \overline{\mathbf{B}(\mathbf{n} \cap^{\psi(y)} \mathbf{n})}.$$

In general there is no simple straightforward description of $\psi(T)$ for a given T . But for $T = (P, Q)$ such that $\forall p \in \langle P \rangle, \forall q \in \langle Q \rangle$ one has $p < q$, the result is very simple.

Lemma. Given tableaux P and Q such that $\forall p \in \langle P \rangle, \forall q \in \langle Q \rangle$ one has $p < q$, then $\psi(P, Q) = (\psi(Q), \psi(P))$. In particular for $T_{\mathcal{I}} = (C_1, \dots, C_l)$ one has $\psi(T_{\mathcal{I}}) = (\psi(C_l), \dots, \psi(C_1))$.

Proof.

This is straightforward from 3.4 and 3.8. Let $P = T(x)$, $Q = T(y)$. Note that for any $a \in \psi(y)$ and any $b \in \psi(x)$ one has $a < b$. Now by 3.4 $(P, Q) = T([x, y])$ and

$$\begin{aligned} \psi(P, Q) &= T(\psi([x, y])) = T([\psi(y), \psi(x)]) && \text{by 3.8} \\ &= (\psi(Q), \psi(P)) && \text{by 3.4.} \end{aligned}$$

■

Note that $\psi(T_{\mathcal{I}}) = (\psi(C_l), \dots, \psi(C_1)) = T_{\psi(\mathcal{I})}$ can be also obtained as a straightforward corollary of the involution ψ of Dynkin diagram.

3.10. As well we can describe explicitly $\psi(T_{\mathcal{I}}(i, \text{next}))$ and $\psi(T_{\mathcal{I}}(i, \text{prev}))$. This requires more subtle consideration of RS insertion. Recall the notation $p(i, \text{next})$ and $p(i, \text{prev})$ from 3.2. Recall the definition $T_{\mathcal{I}}(c_s \swarrow C_i(j))$ from 2.6.

Proposition. *Let $T_{\mathcal{I}} = (C_1, C_2, \dots, C_l)$.*

- (i) *If $T_{\mathcal{I}}(i, \text{next}) \neq \emptyset$ then $\psi(T_{\mathcal{I}}(i, \text{next})) = T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{next})})(c_i))$.*
- (ii) *If $T_{\mathcal{I}}(i, \text{prev}) \neq \emptyset$ then $\psi(T_{\mathcal{I}}(i, \text{prev})) = T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{prev})})(c_{p(i, \text{prev})}))$.*

Proof.

Let us first show (i).

Set $T := T_{\mathcal{I}}(i, \text{next})$. Note that $\text{sh}(T) = \text{sh}(T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{next})})(c_i)))$.

First assume that $i = l$ and $p(l, \text{next}) = 1$. In particular $\text{next}_l(l) = c_1$. Note that for any $\mathcal{I}$ for any i one has $|T^i|$ is the number of chains of length greater or equal than i . In particular by our assumption for $j > 1$ if $c_j > c_l$ then $c_j > c_1$. Thus $|T_{\mathcal{I}}^{c_l+1}| = |T_{\mathcal{I}}^{c_l+2}| = \dots |T_{\mathcal{I}}^{c_1}| =: s$ and $|T_{\mathcal{I}}^{c_l}| = s + 1$, $|T_{\mathcal{I}}^{c_l+1}| = s - 1$. Recall that T is obtained from $T_{\mathcal{I}}$ by moving n from row c_l to row $c_1 + 1$ so that $|T^{c_l}| = |T^{c_l+1}| = \dots = |T^{c_1}| = |T^{c_1+1}| = s$. Let $\{C_1, C_{i_2}, \dots, C_{i_s}\}$ be the chains of length greater than c_l .

Just for the simplicity of further notation note that $C_1 = (1, 2, \dots, c_1)$, $\psi(C_1) = (n + 1 - c_1, \dots, n)$ and $(\psi(C_1))^{c_l+t} = n - c_1 + c_l + t$ for any $t : 0 \leq t \leq c_1 - c_l$

Recall $C_i^{1,j}$ from 2.6. If $c_i \leq j$ then let $C_i^{1,j} = C_i$.

Set $x = [\overline{C_1^{1,c_1}}, \overline{C_2^{1,c_1}}, \dots, \overline{C_{l-1}^{1,c_1}}, \overline{C_l - n}]$, $y = [T^{c_1+1}]$ and $z = w_r(T^{c_1+2, \infty})$. Thus by Part I, 3.2.3 (v) one has $T([z, y, x]) = T$.

Now consider $S := \psi(T) = T(\psi([z, y, x]))$. By 3.8 one has $\psi([y, x, w]) = [\psi(x), \psi(y), \psi(z)]$.

- (1) Put $S' = T([\psi(x)])$ then by 3.9 one has $S' = (\psi(C_l - n), \psi(C_{l-1}^{1,c_1}), \dots, \psi(C_1^{1,c_1}))$. and in particular $\{(n + 1 - T_s^{c_1}) < (n + 1 - T_{s-1}^{c_1}) < \dots < (n + 1 - T_1^{c_1})\} \subset < (S')^1 >$ and $(S')^{c_l} = ((n + 1 - C_{i_s}^{c_1+1-c_l}), \dots, (n + 1 - C_{i_2}^{c_1+1-c_l}), n - c_1 + c_l)$.
- (2) $\psi(y) = [1, n + 1 - T_{s-1}^{c_1+1}, \dots, n + 1 - T_1^{c_1+1}] =: [y_1, \dots, y_s]$. Note that $y_1 < y_2 < \dots < y_s$ and $y_j < (n + 1 - T_{s+1-j}^{c_1})$ for any $j : 1 \leq j \leq s$

- (3) Set $S'' = T([\psi(x), \psi(y)])$. Then by RS procedure and by (1) and (2) one has $(S'')^{c_l+1} = (S')^{c_l}$. In particular $r_{S''}(n - c_1 + c_l) = c_l + 1$.

By RS procedure for any $p \in S''$ one has $r_S(p) \geq r_{S''}(p)$.

In particular $r_S(n - c_1 + c_l) \geq c_l + 1$. Since $S \stackrel{G}{>} T_{\psi(I)}$ we get just by τ -invariant that

$$r_S(n - c_1 + c_l + t) \geq c_l + 1 + t \quad \text{for any } t : 0 \leq t \leq c_1 - c_l. \quad (*)$$

As well one has $\text{sh}(S) = \text{sh}(T)$. The only tableau greater (in geometric order) than $T_{\psi(I)}$ fulfilling $(*)$ and the shape condition is

$$S = T_{\psi(I)}(c_l \swarrow \psi(C_1)(c_l)). \quad (**)$$

Now assume that $i < l$ or $j := p(i, \text{next}) > 1$. Then $T = (P, S, Q)$ where

$$P = (C_1, \dots, C_{j-1}), \quad S = T_{\pi_{\sigma_{j-1}+1, \sigma_i}(I)}(i - j, \text{next}), \quad Q = (C_{i+1}, \dots, C_l).$$

By 3.9 we get $\psi(T) = (\psi(Q), \psi(S), \psi(P))$. By (i) we get that

$$\psi(S) = T_{\psi(\pi_{\sigma_{j-1}+1, \sigma_i}(I))}(c_i \swarrow \psi(C_j)(c_i)).$$

Thus $\psi(T) = T_{\psi(I)}(c_i \swarrow \psi(C_j)(c_i))$. To show (ii) we set $T := T_I(i, \text{prev})$ and again begin

with the case $i = l$ and $p(l, \text{prev}) = 1$. In that case $T = T_I(c_l \swarrow C_1(c_l))$. Note that this case is dual to (i), that is T is S from $(**)$. Thus since ψ is involution we get

$$\psi(T) = T_{\psi(I)}(l, \text{next}) = T_{\psi(I)}(c_l \swarrow \psi(C_1)(c_l)).$$

We proceed as in part (ii) to obtain the result for any i and $p(i, \text{prev})$. ■

3.11. As we have mentioned already in 2.8 if $\text{next}_l(i) = c_i$ then $\text{sh}(T_I(i, \text{next}))$ is obtained from $\text{sh}(T_I)$ by moving a box from row c_i to row $c_i + 1$ so that $\text{codim}_{\mathfrak{m}_I}(\mathcal{V}_{T_I(i, \text{next})}) = 1$. Thus in this case $T_I(i, \text{next})$ is a (geometric) descendant of T_I just by dimension consideration.

To complete the proof of the theorem it remains to show

Proposition. (i) If $\text{next}_l(i) > c_i$ then $T_I(i, \text{next}) \in \mathcal{D}(T_I)$ iff $\text{snext}_r(i) > \text{next}_l(i)$.
(ii) If $\text{prev}_l(i) = \text{sprev}_l(i) \neq n$ then $T_I(i, \text{prev}) \in \mathcal{D}(T_I)$ iff $\text{sprev}_r(i) < \text{sprev}_l(i)$.

Proof.

We begin with (i). If $T_I(i, \text{next})$ is not a descendant, that is if there exists $T : T_I \stackrel{G}{>} T \stackrel{G}{>} T_I(i, \text{next})$, then by 3.5 $T \in T_I(j)$ for some j . Moreover by 3.6 (i) $j > i$. Indeed consider $\pi_{1, C_i^{c_i}}(T)$, $\pi_{1, C_i^{c_i}}(T_I(i, \text{next}))$. Since $\pi_{1, C_i^{c_i}}(T) \neq \pi_{1, C_i^{c_i}}(T_I(i, \text{next}))$ one has by 3.6 (i) that $\pi_{1, C_i^{c_i}}(T) = \pi_{1, C_i^{c_i}}(T_I)$. Thus $j > i$.

Set $\text{sh}(T_{\mathcal{I}}) := \lambda = (\lambda_1, \dots, \lambda_k)$ then $\text{sh}(T_{\mathcal{I}}(i, \text{next})) = \mu$ where

$$\mu_s = \begin{cases} \lambda_s, & \text{if } s \neq c_i, c_{p(i, \text{next})} + 1, \\ \lambda_{c_i} - 1, & \text{if } s = c_i, \\ \lambda_{c_{p(i, \text{next})} + 1} + 1, & \text{if } s = c_{p(i, \text{next})} + 1. \end{cases}$$

Let us first show that if $\text{snext}_r(i) > \text{next}_l(i)$ then $T_{\mathcal{I}}(i, \text{next}) \in \mathcal{D}(T_{\mathcal{I}})$. Assume that $\text{snext}_r(i) > \text{next}_l(i)$. That means that $\forall j : j > i$ one has $c_j \leq c_i$ or $c_j > c_{p(i, \text{next})}$. Then for any $j > i$ if $T_{\mathcal{I}}(j, \text{next})$ is defined then $\text{sh}(T_{\mathcal{I}}(j, \text{next})) = \nu$ where

$$\nu_s = \begin{cases} \lambda_s, & \text{if } s \neq c_j, c_{p(j, \text{next})} + 1; \\ \lambda_{c_j} - 1, & \text{if } s = c_j; \\ \lambda_{c_{p(j, \text{next})} + 1} + 1, & \text{if } s = c_{p(j, \text{next})} + 1. \end{cases}$$

- (1) If $c_j < c_i$ or $c_j > c_{p(i, \text{next})}$ then $\nu \not\prec \mu$.
- (2) If $c_j = c_i$ put $s = C_i^1$, $t = \sigma_j$ and consider $\pi_{s, t}$. Using consequently 3.2 we get

$$\pi_{s, t}(T_{\mathcal{I}}(i, \text{next})) = T_{\pi_{s, t}(\mathcal{I})}, \quad \pi_{s, t}(T_{\mathcal{I}}(j, \text{next})) = T_{\pi_{s, t}(\mathcal{I})}(j - i + 1, \text{next})$$

Hence in both cases $T_{\mathcal{I}}(i, \text{next}) \stackrel{\text{G}}{\not\prec} T_{\mathcal{I}}(j, \text{next})$.

Again, for any $j > i$ if $T_{\mathcal{I}}(j, \text{prev})$ is defined then $\text{sh}(T_{\mathcal{I}}(j, \text{prev})) = \nu$ where

$$\nu_s = \begin{cases} \lambda_s, & \text{if } s \neq c_j + 1, c_{p(j, \text{prev})}; \\ \lambda_{c_{p(j, \text{prev})}} - 1, & \text{if } s = c_{p(j, \text{prev})}; \\ \lambda_{c_j + 1} + 1, & \text{if } s = c_j + 1. \end{cases}$$

and we are left only with the cases when $c_j \neq c_i$ (if $c_j = c_i$ then $T_{\mathcal{I}}(j, \text{prev}) = \emptyset$) so that $\nu \not\prec \mu$.

To complete (i) we must show that if $\text{snext}_r(i) \leq \text{next}_l(i)$ then $T_{\mathcal{I}}(i, \text{next}) \notin \mathcal{D}(T_{\mathcal{I}})$. We use the involution ψ . Consider $T_{\psi(\mathcal{I})}$. Since ψ is order preserving one has that $T_{\mathcal{I}}(i, \text{next}) \in \mathcal{D}(T_{\mathcal{I}})$ iff $\psi(T_{\mathcal{I}}(i, \text{next})) \in \mathcal{D}(T_{\psi(\mathcal{I})})$. Let us show that the last assertion is not true in our case. Indeed, by 3.10

$$\psi(T_{\mathcal{I}}(i, \text{next})) = T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{next})})(c_i)).$$

By 3.5

$$T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{next})})(c_i)) \stackrel{\text{G}}{\geq} T_{\psi(\mathcal{I})}[m]$$

where $m = \psi_{n+1}(C_{p(i, \text{next})}^{c_i})$. Note the the number of the chain $\psi(C_{p(i, \text{next})})$ in $T_{\psi(\mathcal{I})}$ is $l + 1 - p(i, \text{next})$. Since $\text{snext}_r(i) \leq \text{next}_l(i)$ we get that $\text{prev}_l(l + 1 - p(i, \text{next})) > c_i$ in $T_{\psi(\mathcal{I})}$. One has $T_{\psi(\mathcal{I})}[m] = T_{\psi(\mathcal{I})}(l + 1 - p(i, \text{next}), \text{prev}) \neq T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{next})})(c_i))$. Thus by 3.5 $T_{\psi(\mathcal{I})}(c_i \swarrow \psi(C_{p(i, \text{next})})(c_i)) \stackrel{\text{G}}{>} T_{\psi(\mathcal{I})}[m]$ which implies $T_{\psi(\mathcal{I})}(l + 1 - p(i, \text{next}), c_i, c_i) \notin \mathcal{D}(T_{\psi(\mathcal{I})})$. (ii) can be obtained in the same manner as (i) or by applying ψ to (i). We

will show the second way which shows as well that $T_{\mathcal{I}}(i, \text{next}) \in \mathcal{D}(T_{\mathcal{I}})$ if and only if $T_{\psi(\mathcal{I})}(l+1-p(i, \text{next}), \text{prev}) \in \mathcal{D}(T_{\psi(\mathcal{I})})$.

Put $\psi(T_{\mathcal{I}}) := (C'_1, \dots, C'_l)$. Note that $\psi(C_s) = C'_{l+1-s}$ and $|C'_{l+1-s}| = c_s$ for any $s : 1 \leq s \leq l$. Put $s' := l+1-s$ so that $\psi(C_s) = C'_{s'}$. Now set

$$\begin{aligned} \text{snext}_l(s) &:= \begin{cases} \min\{c_j \mid j < s, c_j > c_s\}, & \text{if there exists } j < s \text{ such that } c_j \geq c_s; \\ n, & \text{otherwise.} \end{cases} \\ \text{next}_r(s) &:= \begin{cases} \min\{c_j \mid j > s, c_j \geq c_s\}, & \text{if there exists } j > s \text{ such that } c_j > c_s; \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

Note that for any $s : 1 \leq s \leq l$

- (a) $\text{snext}_r(s')|_{\psi(\mathcal{I})} = \text{snext}_l(s)|_{\mathcal{I}}$ and $\text{next}_l(s')|_{\psi(\mathcal{I})} = \text{next}_r(s)|_{\mathcal{I}}$.
- (b) Assume that s is such that $\text{next}_l(s')|_{\psi(\mathcal{I})} \neq c_s, 0$. Set $p' := p(s', \text{next})|_{\psi(\mathcal{I})}$. Respectively we get $p = l+1-p'$.
 - (b1) If $\text{sprev}_l(p')|_{\psi(\mathcal{I})} \neq n$ then $\text{sprev}_l(p')|_{\psi(\mathcal{I})} = \text{sprev}_r(p)|_{\mathcal{I}}$ and $\text{sprev}_l(p')|_{\psi(\mathcal{I})} < c_s$ (otherwise $\text{next}_l(s')|_{\psi(\mathcal{I})} < c_p$). If $\text{sprev}_l(p')|_{\psi(\mathcal{I})} = n$ then $\text{sprev}_r(p)|_{\mathcal{I}} = 0$. In both cases we get $\text{sprev}_r(p)|_{\mathcal{I}} < c_s$.
 - (b2) One has $\text{prev}_l(p)|_{\mathcal{I}} = \text{sprev}_l(p)|_{\mathcal{I}} = c_s$ iff $\text{snext}_l(s)|_{\mathcal{I}} > \text{next}_r(s)|_{\mathcal{I}}$.

Consider $T_{\mathcal{I}}(i, \text{prev}) \neq \emptyset$. Set $j = p(i, \text{prev})$.

First assume that $T_{\mathcal{I}}(i, \text{prev}) \in \mathcal{D}(T_{\mathcal{I}})$. In that case one has $\psi(T_{\mathcal{I}}(i, \text{prev})) \in \mathcal{D}(T_{\psi(\mathcal{I})})$. By 3.10 (ii) one has $\psi(T_{\mathcal{I}}(i, \text{prev})) = T_{\psi(\mathcal{I})}(c_i \swarrow C'_{j'}, c_j)$. By our assumption one necessarily has by 3.5 that $\psi(T_{\mathcal{I}}(i, \text{prev})) = T_{\psi(\mathcal{I})}(C'_{j'}, \text{next})$ and that $\text{next}_l(j')|_{\psi(\mathcal{I})} = c_i > c_j$. Hence the assumptions of (b) are satisfied for $s = j$. Note that $p(j', \text{next})|_{\psi(\mathcal{I})} = i'$ (otherwise $\text{prev}_l(i)|_{\mathcal{I}} \neq c_j$). Using (b1) to $s = j$ and $p = i$ we get that

$$\text{sprev}_r(i)|_{\mathcal{I}} < c_j. \quad (*)$$

By (i) one has $T_{\psi(\mathcal{I})}(C'_{j'}, \text{next}) \in \mathcal{D}(T_{\psi(\mathcal{I})})$ iff $\text{snext}_r(j')|_{\psi(\mathcal{I})} > \text{next}_l(j')|_{\psi(\mathcal{I})}$. By (a) this is equivalent to $\text{snext}_l(j)|_{\mathcal{I}} > \text{next}_r(j)|_{\mathcal{I}}$. By (b2) this provides $\text{sprev}_l(i)|_{\mathcal{I}} = c_j$. Comparing this to (*) we get $\text{sprev}_r(i)|_{\mathcal{I}} < \text{sprev}_l(i)|_{\mathcal{I}}$.

Now assume that $\text{sprev}_r(i)|_{\mathcal{I}} < \text{sprev}_l(i)|_{\mathcal{I}}$. Recall that $j = p(i, \text{prev})|_{\mathcal{I}}$ that is $c_j = \text{sprev}_l(i)|_{\mathcal{I}}$ and for any $k : j < k < i$ one has that $c_k > c_i$ or $c_k < c_j$. From our assumption one has $\text{next}_r(j)|_{\mathcal{I}} = \text{snext}_r(j)|_{\mathcal{I}} = c_i$ and $\text{snext}_l(j)|_{\mathcal{I}} > c_i$. Thus one has $p(j', \text{next})|_{\psi(\mathcal{I})} = i'$ and $\text{next}_l(j')|_{\psi(\mathcal{I})} < \text{snext}_r(j')|_{\psi(\mathcal{I})}$. Thus by (i) $T_{\psi(\mathcal{I})}(j', \text{next}) \in \mathcal{D}(T_{\psi(\mathcal{I})})$. Applying ψ we get $\psi(T_{\psi(\mathcal{I})}(j', \text{next})) \in \mathcal{D}(T_{\mathcal{I}})$. On the other hand by 3.10 $\psi(T_{\psi(\mathcal{I})}(j', \text{next})) = T_{\mathcal{I}}(c_j \swarrow C'_i(c_j)) = T_{\mathcal{I}}(C_i, \text{prev})$. Hence $T_{\mathcal{I}}(C_i, \text{prev}) \in \mathcal{D}(T_{\mathcal{I}})$. ■

4. On geometry of descendants of a Richardson orbital variety

4.1. As it was shown in [Me] one has $\overline{\mathbf{n} \cap \mathcal{O}} = \mathbf{n} \cap \overline{\mathcal{O}}$ for any $\mathcal{O}$. The question is whether considering $\mathbf{m}_{\mathcal{I}}$ instead of $\mathbf{n}$ gives the same equality. Of course the first restriction is $\mathcal{O} \geq \mathcal{O}_{\mathcal{I}}$ since if $\mathcal{O} < \mathcal{O}_{\mathcal{I}}$ then $\overline{\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}} = \emptyset$ and $\mathbf{m}_{\mathcal{I}} \cap \overline{\mathcal{O}} = \mathbf{m}_{\mathcal{I}}$ so the equality trivially fails to be true. Let us show that even for $\mathcal{O} > \mathcal{O}_{\mathcal{I}}$ the equality does not necessarily hold. Recall that

$$\mathbf{m}_{\mathcal{I}} \cap \overline{\mathcal{O}} = \coprod_{\mathcal{O}' \geq \mathcal{O}} \mathbf{m}_{\mathcal{I}} \cap \mathcal{O}'.$$

Now take $\mathcal{I} \subset \Pi$ for which there exist $\mathcal{V} \in \mathcal{D}_G(\mathcal{V}_{\mathcal{I}})$ such that $\mathcal{O}_{\mathcal{V}}$ is not a descendant of $\mathcal{O}_{\mathcal{I}}$. Let $\mathcal{O}$ be an intermediate orbit, that is $\mathcal{O}_{\mathcal{I}} < \mathcal{O} < \mathcal{O}_{\mathcal{V}}$. Obviously $\mathcal{V} \subset \mathbf{m}_{\mathcal{I}} \cap \overline{\mathcal{O}}$. On the other hand one has by 2.3

$$\mathbf{m}_{\mathcal{I}} \cap \mathcal{O} = \cup \mathcal{V}_i,$$

where $\mathcal{V}_i$ are orbital varieties. Thus $\overline{\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}} = \cup \overline{\mathcal{V}_i}$. Now by 2.8 one has $\mathcal{V} \not\subset \overline{\mathcal{V}_i}$ for any i thus $\mathcal{V} \not\subset \cup \overline{\mathcal{V}_i}$ just by irreducibility of $\mathcal{V}$.

The first example occurs in $\mathfrak{sl}_4$. Let

$$T_{\mathcal{I}} = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & & \\ \hline \end{array}, \quad S = \begin{array}{|c|c|} \hline 1 & 3 \\ \hline 2 & 4 \\ \hline \end{array} \quad \text{and} \quad T = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & \\ \hline 3 & \\ \hline \end{array}.$$

Take $\lambda = (2, 2)$. One has $\mathcal{V}_T \subset \mathbf{m}_{\mathcal{I}} \cap \overline{\mathcal{O}_{\lambda}}$ and $\mathcal{V}_T \not\subset \overline{\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}_{\lambda}} = \overline{\mathcal{V}_S}$.

On the other hand note that by 2.3 and the definition of $\mathcal{D}_G(T_{\mathcal{I}})$ one has

$$\overline{\mathbf{m}_{\mathcal{I}} \cap \left(\bigcup_{T \in \mathcal{D}_G(T_{\mathcal{I}})} \mathcal{O}_{\text{sh } T} \right)} = \mathbf{m}_{\mathcal{I}} \cap \left(\bigcup_{T \in \mathcal{D}_G(T_{\mathcal{I}})} \overline{\mathcal{O}_{\text{sh } T}} \right)$$

In particular if $|T_{\mathcal{I}}^1| = 2$ or $|T_{\mathcal{I}}^1| = |T_{\mathcal{I}}^2| = \dots = |T_{\mathcal{I}}^k|$ that is in the cases when all $T \in \mathcal{D}_G(T_{\mathcal{I}})$ are of the same shape, call it λ , one has $\overline{\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}_{\lambda}} = \mathbf{m}_{\mathcal{I}} \cap \overline{\mathcal{O}_{\lambda}}$.

4.2. As we have explained in 1.11 all the descendants of codimension 1 of $\mathcal{V}_{\mathcal{I}}$ are complete intersections. According to 2.8 some of the descendants of $\mathcal{V}_{\mathcal{I}}$ are of codimension greater than 1 in it. The question is whether a descendant of codimension greater than 1 of $\mathcal{V}_{\mathcal{I}}$ is necessarily a complete intersection. In 4.4 we give an example in $\mathfrak{sl}_6$ of $\mathcal{V}_{\mathcal{I}}$ and its descendant of codimension 2 which is not a complete intersection.

First of all note that for $\mathfrak{sl}_n$ where $n \leq 5$ all the descendants of a Richardson orbital variety are complete intersections. Indeed as it is noted in [vanL, p.16] all the orbital varieties for $n \leq 4$ are complete intersections and the only orbital variety in $\mathfrak{sl}_5$ which is

not a complete intersection is $\mathcal{V}_T$ where

$$T = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline 5 & \\ \hline \end{array}$$

Note that $\tau(T) = \{\alpha_2, \alpha_4\} =: \mathcal{I}$ and that by 2.8 $T \notin \mathcal{D}_G(T_{\mathcal{I}})$.

4.3. Before considering the example in $\mathfrak{sl}_6$ let us introduce the notation we use in what follows.

Let $e_{i,j}$ be the matrix having 1 in the ij -th entry and 0 elsewhere. Let $x_{j,i}$ denote the coordinate function on $\mathfrak{g} = \mathfrak{sl}_n$ defined by

$$x_{j,i}(e_{r,s}) = \begin{cases} -1, & \text{if } (r,s) = (i,j), \\ 0, & \text{otherwise.} \end{cases}$$

Then the Poisson bracket $\{, \}$ defined on $S(\mathfrak{g}^*)$ through the Lie bracket on $\mathfrak{g}$ satisfies

$$\{x_{i,j}, x_{r,s}\} = \delta_{j,r}x_{i,s} - \delta_{s,i}x_{r,j} \quad (*)$$

where $\delta_{i,j}$ is the Kronecker delta. Let $\mathfrak{m}^-$ be the opposed algebra of $\mathfrak{m}$. Setting $x_{i,j} = e_{i,j}$ for $i > j$ identifies $\mathfrak{m}^-$ with $\mathfrak{m}^*$. Set $A := S(\mathfrak{m}^-)$ that is a symmetric algebra of $\mathfrak{m}^-$.

4.4. Consider $\mathfrak{sl}_6$. Set $\mathcal{I} = \{\alpha_1, \alpha_2, \alpha_5\}$. One has

$$T_{\mathcal{I}} = \begin{array}{|c|c|c|} \hline 1 & 4 & 5 \\ \hline 2 & 6 & \\ \hline 3 & & \\ \hline \end{array}$$

Consider $\mathcal{V} := \mathcal{V}_T$ where

$$T = \begin{array}{|c|c|c|} \hline 1 & 4 & 5 \\ \hline 2 & & \\ \hline 3 & & \\ \hline 6 & & \\ \hline \end{array}$$

Note that $T \in \mathcal{D}_G(T_{\mathcal{I}})$ and moreover $\text{sh}(T) = (3, 1, 1, 1)$ is a descendant of $\text{sh}(T_{\mathcal{I}}) = (3, 2, 1)$. One has $\text{codim}_{\mathfrak{m}_{\mathcal{I}}}(\mathcal{V}) = 2$.

Let $I(\mathcal{V})$ denote the ideal of definition of $\mathcal{V}$ in A . Let us show that $I(\mathcal{V}) = \sqrt{I}$ where $I := \langle f_1, f_2, f_3, f_4 \rangle$ for

$$\begin{aligned}
f_1 &= \begin{vmatrix} x_{4,2} & x_{4,3} & 0 \\ x_{5,2} & x_{5,3} & x_{5,4} \\ x_{6,2} & x_{6,3} & x_{6,4} \end{vmatrix} = x_{4,2}x_{5,3}x_{6,4} + x_{4,3}x_{5,4}x_{6,2} - x_{4,2}x_{5,4}x_{6,3} - x_{4,3}x_{5,2}x_{6,4} \\
f_2 &= \begin{vmatrix} x_{4,1} & x_{4,3} & 0 \\ x_{5,1} & x_{5,3} & x_{5,4} \\ x_{6,1} & x_{6,3} & x_{6,4} \end{vmatrix} = x_{4,1}x_{5,3}x_{6,4} + x_{4,3}x_{5,4}x_{6,1} - x_{4,1}x_{5,4}x_{6,3} - x_{4,3}x_{5,1}x_{6,4} \\
f_3 &= \begin{vmatrix} x_{4,1} & x_{4,2} & 0 \\ x_{5,1} & x_{5,2} & x_{5,4} \\ x_{6,1} & x_{6,2} & x_{6,4} \end{vmatrix} = x_{4,1}x_{5,2}x_{6,4} + x_{4,2}x_{5,4}x_{6,1} - x_{4,1}x_{5,4}x_{6,2} - x_{4,2}x_{5,1}x_{6,4} \\
f_4 &= \begin{vmatrix} x_{4,1} & x_{4,2} & x_{4,3} \\ x_{5,1} & x_{5,2} & x_{5,3} \\ x_{6,1} & x_{6,2} & x_{6,3} \end{vmatrix} = \\
&= x_{4,1}x_{5,2}x_{6,3} + x_{4,2}x_{5,3}x_{6,1} + x_{4,3}x_{5,1}x_{6,2} - x_{4,1}x_{5,3}x_{6,2} - x_{4,2}x_{5,1}x_{6,3} - x_{4,3}x_{5,2}x_{6,1}
\end{aligned}$$

Let us show first of all that all four polynomials are irreducible. Indeed $f_1 = m_{l_{\mathcal{I}'}}$ where $\mathbf{m}_{\mathcal{I}'} = \pi_{2,6}(\mathbf{m}_{\mathcal{I}})$. Thus f_1 is irreducible by [JM]. Now $f_2 = -\{x_{2,1}, f_1\}$ and $f_3 = \{x_{3,1}, f_1\}$ so they are irreducible. Finally $f_4 = m_{l_{\mathcal{I}''}}$ where $\mathcal{I}'' = \{\alpha_1, \alpha_2, \alpha_4, \alpha_5\}$ and again is irreducible by [JM]. Further note that $f_1 \notin \langle f_2, f_3, f_4 \rangle$ since $f_2|_{x_{4,1}=x_{5,1}=x_{6,1}=0} = f_3|_{x_{4,1}=x_{5,1}=x_{6,1}=0} = f_4|_{x_{4,1}=x_{5,1}=x_{6,1}=0} = 0$ and $f_1|_{x_{4,1}=x_{5,1}=x_{6,1}=0} \neq 0$. Exactly in the same way we show that $f_2 \notin \langle f_1, f_3, f_4 \rangle$, $f_3 \notin \langle f_1, f_2, f_4 \rangle$ and $f_4 \notin \langle f_1, f_2, f_3 \rangle$. Thus I is generated by 4 polynomials. One can see at once that I is stable under Poisson bracket action.

Let $\mathcal{V}(I)$ denote variety of I . It consists of all the matrices in $\mathbf{m}_{\mathcal{I}}$ of rank less or equal to 2 that is $\mathcal{V}(I) = \mathbf{m} \cap (\mathcal{O}_{\lambda_1} \cup \mathcal{O}_{\lambda_2} \cup \mathcal{O}_{\lambda_3} \cup \mathcal{O}_{\lambda_4})$ where $\lambda_1 = (3, 1, 1, 1)$, $\lambda_2 = (2, 2, 1, 1)$, $\lambda_3 = (2, 1, 1, 1, 1)$, $\lambda_4 = (1, 1, 1, 1, 1, 1)$. In particular $\overline{\mathcal{V}}_T \subset \mathcal{V}(I)$. By 2.3 one has

$$\begin{aligned}
\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}_{\lambda_1} &= \mathcal{V}_T \\
\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}_{\lambda_2} &= \mathcal{V}_P \cup \mathcal{V}_Q \\
\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}_{\lambda_3} &= \mathcal{V}_S \cup \mathcal{V}_U \\
\mathbf{m}_{\mathcal{I}} \cap \mathcal{O}_{\lambda_4} &= \mathcal{V}_Y
\end{aligned}$$

where

$$P = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & \\ \hline 6 & \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|} \hline 1 & 5 \\ \hline 2 & 6 \\ \hline 3 & \\ \hline 4 & \\ \hline \end{array}, \quad S = \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & \\ \hline 3 & \\ \hline 5 & \\ \hline 6 & \\ \hline \end{array}, \quad U = \begin{array}{|c|c|} \hline 1 & 5 \\ \hline 2 & \\ \hline 3 & \\ \hline 4 & \\ \hline 6 & \\ \hline \end{array} \quad \text{and} \quad Y = \begin{array}{|c|} \hline 1 \\ \hline 2 \\ \hline 3 \\ \hline 4 \\ \hline 5 \\ \hline 6 \\ \hline \end{array}.$$

By Part I, 3.3.3 one can see that $P, Q, S, U, Y \stackrel{D}{>} T$. Thus $\mathcal{V}(I) \subset \overline{\mathcal{V}}_T$. We get $\mathcal{V}(I) = \overline{\mathcal{V}}_T$.

To show that $\overline{\mathcal{V}}_T$ is not a complete intersection we must show that $\sqrt{I}$ is generated by at least 3 polynomials. We will show that it is generated by at least 4 polynomials.

Let $\sqrt{I} = \langle p_i \rangle_{i=1}^k$. Let us write $p_i = \sum p_i^j$ where p_i^j is a component of p_i whose monomials contain j different variables.

First let us show that $p_i^j = 0$ for $j \leq 2$.

- (i) It is obvious that $p_i^0 = 0$ since there exists t_i such that $p_i^{t_i} \in I$.
- (ii) Note that A has 11 variables $x_{i,j}$. Note that for any specialization $\overline{x}$ of $\{x_{i,j}\}$ taking 10 variables to 0 one has $f_i|_{\overline{x}} = 0$ for $i = 1, 2, 3, 4$. Thus $I|_{\overline{x}} = 0$. On the other hand $p_i|_{\overline{x}} = p_i^1|_{\overline{x}}$ and $p_i^{t_i}|_{\overline{x}} = (p_i^1)^{t_i}|_{\overline{x}} \in I|_{\overline{x}}$. Thus $p_i^1|_{\overline{x}} = 0$ for any such specialization which provides $p_i^1 = 0$.
- (iii) Now note that for any specialization $\overline{x}$ of $\{x_{i,j}\}$ taking 9 variables to 0 one has $f_i|_{\overline{x}} = 0$ for $i = 1, 2, 3, 4$. Thus exactly in the same way as in (ii) we get that $p_i^2 = 0$.

Now let $p_{i,j}$ denote the component of p_i of total degree j . Then $p_i = p_{i,3} + \hat{p}_i$ where $\hat{p}_i = \sum_{j \geq 4} p_{i,j}$. Since $f_1, f_2, f_3, f_4 \in \sqrt{I}$ we get by degree considerations

$$f_j = \sum_i B_i^j p_i = \sum_i b_i^j p_{i,3}$$

where B_i^j are some polynomials and $b_i^j \in \mathbb{C}$. Consider a linear affine space $V = \mathbb{C}^r$ with the basis $\{x_{i,j}x_{p,q}x_{s,t}\}$ where $(i,j), (p,q), (s,t)$ are pairwise different and $r = \binom{11}{3}$. Since $f_1, \dots, f_4$ are linearly independent in V one has that the number of p_i must be at least 4.

Moreover using more subtle analysis one can show that for any $i : 1 \leq i \leq k$ and any $j : j \geq 3$ one has $p_{i,j} \in \sqrt{I}$ and then using specializations taking $\sqrt{I}$ onto $I_i = \langle f_i \rangle$ which are all simple ideals and some further specializations one can show that $\sqrt{I} = \langle f_1, f_2, f_3, f_4 \rangle$.

INDEX OF NOTATION

- | | |
|--|---|
| 1.2 $\mathcal{O}, \mathcal{V}, \mathcal{O}_{\mathcal{V}}$ | 2.2 $T_j^i, \omega^i(T), T_j, T^i, T^{i,j}, T^{i,\infty}, (T_j) ,$ |
| 1.3 $R, R^+, \Pi, X_\alpha, \mathbf{B}, \mathfrak{b}, \mathcal{V}_w, \mathcal{O}_w$ | $ (T^i) , <T>, \mathbb{T}_n, \mathbb{T}_E, (T, S), T^\dagger$ |
| 1.4 $\mathcal{I}, \mathbf{P}_{\mathcal{I}}, \mathbf{M}_{\mathcal{I}}, \mathbf{m}_{\mathcal{I}},$ | 2.4 $T_{\mathcal{I}}, C_i^{\mathcal{I}}, c_i, \sigma_i$ |
| $W_{\mathcal{I}}, w_{\mathcal{I}}, \mathcal{V}_{\mathcal{I}}, \mathcal{O}_{\mathcal{I}}$ | 2.6 $\hat{T}_{s,t}, C_i^{1,j-1}, C_i(j), T_{\mathcal{I}}(c_{s \setminus} C_i(j))$ |
| 1.5 $\lambda, \mathcal{O}_\lambda, \lambda \leq \mu, \mathcal{O} \leq \mathcal{O}'$ | 2.7 $\text{next}_l(i), \text{prev}_l(i), \text{sprev}_l(i), \text{snext}_r(i),$ |
| 1.6 $\mathcal{C}_{\mathcal{V}}, \mathcal{C}_w, \mathbf{S}_n, T(w),$ | $\text{sprev}_r(i), T_{\mathcal{I}}(i, \text{next}), T_{\mathcal{I}}(i, \text{prev}), T_{\mathcal{I}}(i)$ |
| $\mathbf{T}_n, \mathcal{V}_T, T_{\mathcal{V}}, \text{sh}(T)$ | 2.8 $\mathcal{D}_G(T)$ |
| 1.7 $r_T(u), \tau(T), \tau(\mathcal{V})$ | 3.1 $<i, j>, \pi_{i,j}$ |
| 1.8 $\mathcal{V} \stackrel{\text{G}}{\leq} \mathcal{W}$ | 3.2 $\mathcal{I}_n, \mathcal{I}_1, p(i, \text{next}), p(i, \text{prev})$ |
| 2.1 $S(w), y \stackrel{\text{D}}{\leq} w, \tau(w), \mathbf{P}_\alpha, \alpha_{i,j},$ | 3.4 $\binom{P}{Q}$ |
| $p_w(i), <w>, \overline{w}, [x, y], \mathbb{S}_n, \mathbb{S}_E$ | 4.3 $x_{i,j}, \mathbf{m}^-, A$ |

REFERENCES

- [1] A. Joseph, On the variety of a highest weight module, J. of Algebra 88 (1984), 238-278.
- [2] A. Joseph, A. Melnikov, Quantization of hypersurface orbital varieties in $\mathfrak{sl}_n$, The orbit method in geometry and physics. In honor of A.A. Kirillov, series "Progress in Mathematics", 213, Birkhauser, 2003, 165-196.
- [3] A. Melnikov, On orbital variety closures in $\mathfrak{sl}_n$. I. Induced Duflo order, J. of Algebra, to appear.
- [4] D. Mertens, Über einen satz von Spaltenstein, preprint.
- [5] N. Spaltenstein, On the fixed point set of a unipotent element on the variety of Borel subgroups, Topology 16 (1977), 203-204.
- [6] N. Spaltenstein, The fixed point set of a unipotent transformation on the flag manifold, Proc. Konin. Nederl. Akad. 79 (1976), 452-456.
- [7] M. A. van Leeuwen, The Robinson-Schensted and Schützenberger algorithms, Part II: Geometric interpretations, CWI report AM-R9209 (1992).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA 31905, ISRAEL

E-mail address: melnikov@math.haifa.ac.il